

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/001624

International filing date: 17 February 2005 (17.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04003554.5
Filing date: 17 February 2004 (17.02.2004)

Date of receipt at the International Bureau: 29 April 2005 (29.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

PCT/EP05/09624

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04003554.5

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

100-100000

100-100000

100-100000

100-100000



Anmeldung Nr:
Application no.: 04003554.5
Demande no:

Anmeldetag:
Date of filing: 17.02.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

National Public Health Institute
Mannerheimintie 166
00300 Helsinki
FINLANDE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Identification of SNPs associated with hyperlipidemia, dyslipidemia and defective
carbohydrate metabolism

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

17 Feb. 2004

5 New EP Application
National Public Health Institute et al.
Our Ref.: K 1114EP

10 **IDENTIFICATION OF SNPs ASSOCIATED WITH HYPERLIPIDEMIA,
DYSLIPIDEMIA AND DEFECTIVE CARBOHYDRATE METABOLISM**

The present invention relates to a nucleic acid molecule comprising a chromosomal region contributing to or indicative of hyperlipidemias and/or dyslipidemias and/or defective carbohydrate metabolism, wherein said nucleic acid molecule is selected
15 from the group consisting of: (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence has one or more mutations having an effect on USF1 function; (b) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence is characterized by comprising a guanine or an adenine
20 residue in position 3966 in intron 7 of the USF1 sequence; and/or (c) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence is characterized by comprising a cytosine or a thymine residue in position 5205 in exon 11 of the USF1 sequence; wherein said nucleic molecule extends, at a maximum, 50000 nucleotides over the 5' and/or 3' end of the
25 nucleic acid molecule of SEQ ID NO: 1. The present invention further relates to a diagnostic composition comprising a nucleic acid molecule encoding USF1 or a fragment thereof, the nucleic acid molecule disclosed herein, the vector, the primer or primer pair of the present invention or an antibody specific for USF1. Finally, the present invention relates to the use of the nucleic acid molecule of the invention for
30 the preparation of a pharmaceutical composition for the treatment of hyperlipidemia, dyslipidemia, coronary heart disease, type II diabetes, metabolic syndrome, hypertension or atherosclerosis.

A variety of documents is cited throughout this specification. The disclosure content of these documents, including manufacturer's manuals and catalogues, is herewith
35 incorporated by reference.

Familial combined hyperlipidemia (FCHL) is characterized by elevated levels of serum total cholesterol (TC), triglycerides (TG), or both^{1,2}. Recently, the first major

5 locus for FCHL was identified on human chromosome 1q21-q23 in 31 Finnish FCHL families⁴. This finding has been replicated in FCHL families from other, more heterogeneous populations⁵⁻⁷. In addition, genome-wide scans have identified several other putative loci for FCHL in Finnish and Dutch study samples⁸⁻⁹. Interestingly, the same markers in the 1q21 region have also been linked to type 2
10 diabetes mellitus (T2DM) in numerous studies¹⁰⁻¹⁴, including a Finnish study¹⁵. The evidence for linkage obtained for 1q21 has varied in these FCHL and T2DM studies, most likely reflecting genetic heterogeneity as well as population-based and diagnostic differences. Importantly, however, many of the critical metabolic features of FCHL, e.g. hypertriglyceridemia and insulin resistance, also represent trait
15 components of T2DM. Interestingly, a rodent locus for combined hyperlipidemia was linked to a region on mouse chromosome 3, potentially orthologous with human 1q21 (ref. 16). The underlying gene, thioredoxin interacting protein (*TXNIP*), was recently identified providing a strong positional candidate for human FCHL¹⁷.

As pointed out above, familial combined hyperlipidemia (FCHL) is characterized by
20 elevated levels of serum total cholesterol (TC), triglycerides (TG), or both^{1,2}. This complex disorder is the most common familial hyperlipidemia with a prevalence of 1% to 2% in Western populations¹. FCHL constitutes a powerful genetic factor in atherosclerosis since it is observed in about 20% of coronary heart disease (CHD) patients under 60 years³. Despite tremendous efforts to identify the molecular
25 mechanisms underlying FCHL, its etiology remains unknown. As a consequence it is presently not possible to diagnose or treat patients affected by familial combined hyperlipidemia (FCHL).

In view of the above, the technical problem underlying the present invention was to provide means and methods that allow for an accurate and convenient diagnosis of
30 of hyperlipidemias and/or dyslipidemias or defective carbohydrate metabolism or of a predisposition to these conditions.

The solution to said technical problem is achieved by the embodiments characterized in the claims.

Thus, the present invention relates to a nucleic acid molecule comprising a
35 chromosomal region contributing to or indicative of hyperlipidemias and/or

5 dyslipidemias or defective carbohydrate metabolism, wherein said nucleic acid
molecule is selected from the group consisting of: (a) a nucleic acid molecule having
or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid
sequence has one or more mutations having an effect on USF1 function; (b) a
10 nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID
NO: 1, wherein said nucleic acid sequence is characterized by comprising a guanine
or an adenine residue in position 3966 in intron 7 of the USF1 sequence; and/or (c)
a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID
NO: 1, wherein said nucleic acid sequence is characterized by comprising a
15 cytosine or thymine residue in position 5205 in exon 11 of the USF1 sequence;
wherein said nucleic molecule extends, at a maximum, 50000 nucleotides over the
5' and/or 3' end of the nucleic acid molecule of SEQ ID NO: 1. In preferred
embodiments, the nucleic acid molecule extends up to 40000 nucleotides or up to
25000 nucleotides or up to 5000 nucleotides over the 5' and/or 3' end of the nucleic
acid molecule of SEQ ID NO: 1.

20 The term "hyperlipidemias and dyslipidemias" refers to diseases associated with an
increased levels of serum total cholesterol and/or triglycerides, as well as increased
levels of low-density lipoprotein (LDL) cholesterol and/or apolipoprotein B and/or
decreased levels of serum high-density lipoprotein (HDL) cholesterol and/or small
dense LDL. In accordance with the present invention such diseases include familial
25 combined hyperlipidemia (FCHL), hypercholesterolemia, hypertriglyceridemia,
hypoalphalipoproteinemia, hyperapobetalipoproteinemia (hyperapoB), familial
dyslipidemic hypertension (FDH), hypertension, coronary heart disease and
atherosclerosis.

30 In accordance with the invention, the term "defective carbohydrate metabolism"
refers to glucose intolerance and insulin resistance. Defective carbohydrate
metabolism might therefore be indicative of diseases such as type 2 diabetes
mellitus (T2DM) and metabolic syndrome.

35 The term "contributing to or indicative of hyperlipidemias and/or dyslipidemias or
defective carbohydrate metabolism", refers to the fact that the SNPs and thus the
corresponding nucleic acid molecules found are indicative of the condition and

5 possibly also causative therefore. Accordingly, this term necessarily requires that the recited position is indicative of the condition. Said term, on the other hand, does not necessarily require that the particular position containing the SNP is actually causative or contributes to the condition. Yet, said term does not exclude a causative or contributory role of either or both SNPs.

10 The nucleotide sequence designated SEQ ID NO:1 is a genomic nucleotide sequence of 5687 bp, representing USF1 as deposited under databank accession number RefSeq: NM_007122 for the human USF1 mRNA with the corresponding genomic sequence as deposited under >hg16_refGene_NM_007122
15 range=chr1:158225833-158231519 in the UCSC Genome Browser on Human in July 2003. For the purpose of the present invention, the activity or function of the polypeptide encoded by this nucleotide sequence is defined as "wild-type USF1 protein activity". Likewise, SEQ ID NO:1 is understood as representing wild-type USF1 if sequence position 3966 is an adenine and sequence position 5205 is a thymine. USF1 is known as a transcription factor, capable of binding to the
20 recognition sequence CACGTG termed E box and capable of regulating the expression of genes such as apolipoproteins CIII (APOC3), AII (APOA2), APOE, hormone sensitive lipase (LIPe), fatty acid synthase (FAS), glucokinase (GCK), glucagon receptor (GCGR), ATP-binding cassette, subfamily A (ABCA1), renin (REN) and angiotensinogen (AGT). Moreover, USF1 is known to interact with other
25 factors of the cellular transcription machinery, such as USF2.

The term "one or more mutations having an effect on USF1 function" refers to mutations affecting USF1 function. Throughout the present invention the term "function" and "activity" are used exchangeable. Since USF1 is a transcription factor, the term "USF1 function" refers to its activity as a transcription factor including its
30 specificity to its target recognition sequence on the genomic DNA, its protein interaction sequences and its capability of modulating or regulating transcription. It is important to note, however, that also mutations outside of the coding region of USF1 can have an effect on USF1 function. Such mutations are, for example, mutations affecting the amount of USF1 transcribed in a cell (including mutations
35 affecting promoter activity) or mutations that have an impact on splicing or

- 5 intracellular transport of the RNA transcripts. Any of these mutations is also comprised by the present invention.

The term "nucleic acid molecule" refers both to naturally and non-naturally occurring nucleic acid molecules. Non-naturally occurring nucleic acid molecules include cDNA as well as derivatives such as PNA.

- 10 The term "nucleic acid molecule [...] comprising the nucleic acid sequence of SEQ ID NO:", as used throughout this specification, refers to nucleic acid molecules that are at least 1 nucleotide longer than the nucleic acid molecule specified by the SEQ ID NO. At the same time, these nucleic acid molecules extend, at a maximum, 50000 nucleotides over the 5' and/or 3' end of the nucleic acid molecule of the
15 invention specified e.g. by the SEQ ID NO: 1.

- A number of previous studies in mammalia have tried to identify chromosomal regions contributing to or associated with familial combined hyperlipidemia. A rodent locus for combined hyperlipidemia was linked to a region on mouse chromosome 3, potentially orthologous with human 1q21 (ref. 16). The underlying gene, thioredoxin
20 interacting protein (*TXNIP*), was recently identified providing a strong positional candidate for human FCHL¹⁷. Surprisingly, the results disclosed by the present invention show that two single-nucleotide polymorphisms located in intron 7 and exon 11, respectively, of human USF1 are associated with hyperlipidemias, dyslipidemias and defective carbohydrate metabolism. The disclosed
25 polymorphisms allow to screen individuals for a presence or predisposition of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism.

- In a preferred embodiment, the nucleic acid molecule of the present invention is genomic DNA. This preferred embodiment of the invention reflects the fact that usually the analysis would be carried out on the basis of genomic DNA from body
30 fluid, cells or tissue isolated from the person under investigation. In a further preferred embodiment of the nucleic acid molecule of the invention said genomic DNA is part of a gene. In accordance with the invention, it is preferred that at least intron 7 of the USF1 gene harboring SNP1 in position 3966 and/or exon 11 of the USF1 gene harboring SNP2 in position 5205 relative to the USF1 gene is analyzed.

5 It is a central aspect of the present invention that a guanine residue in position 3966
of the USF1 gene indicates the presence of a disease-associated allele, whereas an
adenine residue in the same position of the USF1 gene is indicative for the healthy
allele. Likewise, a cytosine residue in position 5205 of the USF1 gene indicates the
presence of a disease-associated allele, whereas a thymine residue is indicative for
10 the healthy allele.

The present invention also relates to a fragment of the nucleic acid molecule the
present invention having at least 20 nucleotides wherein said fragment comprises
nucleotide position 3966 and/or position 5205 of SEQ ID NO:1. The fragment of the
invention may be of natural as well as of (semi)synthetic origin. Thus, the fragment
15 may, for example, be a nucleic acid molecule that has been synthesized according
to conventional protocols of organic chemistry. Importantly, the nucleic acid
fragment of the invention comprises nucleotide position 3966 in intron 7 of the USF1
gene or nucleotide position 5205 in exon 11 of the USF1 gene. In these positions,
the fragment may have either the wild-type nucleotide or the nucleotide contributing
20 to or indicative of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate
metabolism (also referred to as the "mutant" or "disease-associated" sequence).
Consequently, the fragment of the invention may be used, for example, in assays
differentiating between the wild-type and the mutant sequence.

It is further preferred that the fragment of the invention consists of at least 17
25 nucleotides, more preferred at least 20 nucleotides, and most preferred at least 25
nucleotides such as 30 nucleotides. Preferably, however, the fragment is of up to
100bp, up to 200bp, up to 300bp, up to 400bp, up to 500bp, up to 600bp, up to
700bp, up to 800bp, up to 900bp or up to 1000bp in length.

Furthermore, the invention relates to a nucleic acid molecule which is
30 complementary to the nucleic acid molecule of the present invention and which has
a length of at least 17 or of at least 20 nucleotides. Preferably, however,
complementary nucleic acid molecule is of up to 100bp, up to 200bp, up to 300bp,
up to 400bp, up to 500bp, up to 600bp, up to 700bp, up to 800bp, up to 900bp or up
to 1000bp in length.

5 This embodiment of the invention comprising at least 15 or at least 20 nucleotides and covering at least position 3966 or position 5205 of the USF1 gene is particularly useful in the analysis of the genetic setup in the recited positions in hybridization assays. Thus, for example, a 15 mer exactly complementary either to the wild-type
10 dyslipidemia and/or defective carbohydrate metabolism may be used to differentiate between the polymorphic variants. This is because a nucleic acid molecule labeled with a detectable label not exactly complementary to the DNA in the analyzed sample will not give rise to a detectable signal, if appropriate hybridization and washing conditions are chosen.

15 In this regard, it is important to note that the nucleic acid molecule of the invention, the fragment thereof as well as the complementary nucleic acid molecule may be detectably labeled. Detectable labels include radioactive labels such as ^3H , or ^{32}P or fluorescent labels. Labeling of nucleic acids is well understood in the art and described, for example, in Sambrook et al., "Molecular Cloning, A Laboratory
20 Manual"; ISBN: 0879695765, CSH Press, Cold Spring Harbor, 2001.

Hybridisation is preferably performed under stringent or highly stringent conditions. "Stringent or highly stringent conditions" of hybridization are well known to or can be established by the person skilled in the art according to conventional protocols. Appropriate stringent conditions for each sequence may be established on the basis
25 of well-known parameters such as temperature, composition of the nucleic acid molecules, salt conditions etc.: see, for example, Sambrook et al., "Molecular Cloning, A Laboratory Manual"; ISBN: 0879695765, CSH Press, Cold Spring Harbor, 2001 and earlier edition Sambrook et al., "Molecular Cloning, A Laboratory Manual"; CSH Press, Cold Spring Harbor, 1989 or Higgins and Hames (eds.),
30 "Nucleic acid hybridization, a practical approach", IRL Press, Oxford 1985 (reference 54), see in particular the chapter "Hybridization Strategy" by Britten & Davidson, 3 to 15. Typical (highly stringent) conditions comprise hybridization at 65°C in 0.5xSSC and 0.1% SDS or hybridization at 42°C in 50% formamide, 4xSSC and 0.1% SDS. Hybridization is usually followed by washing to remove unspecific
35 signal. Washing conditions include conditions such as 65°C , 0.2xSSC and 0.1% SDS or 2xSSC and 0.1% SDS or 0.3xSSC and 0.1% SDS at 25°C – 65°C .

- 5 Hybridisation may also be performed under conditions of lower stringency. The parameters of such hybridization conditions are described in Sambrook et al., "Molecular Cloning, A Laboratory Manual"; ISBN: 0879695765, CSH Press, Cold Spring Harbor, 2001 in more detail. A non-limiting, example of low stringency hybridization conditions are hybridization in 35% formamide, 5.times. SSC, 50 mM
- 10 Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40.degree. C., followed by one or more washes in 2.times. SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50.degree. C. Other conditions of low stringency that may be used are well known in the art (e.g., as employed for cross-species
- 15 hybridizations). See, e.g., Ausubel, et al. (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons. NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY; Shilo and Weinberg, 1981, Proc Natl Acad Sci USA 78: 6789-6792.

- 20 In addition, the invention relates to a vector comprising the nucleic acid molecule as described herein above. The vectors may particularly be plasmids, cosmids, viruses or bacteriophages used conventionally in genetic engineering that comprise the nucleic acid molecule of the invention. Preferably, said vector is an expression vector and/or a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses,
- 25 or bovine papilloma virus, may be used for delivery of the nucleic acid molecule of the invention into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook et al., loc. cit. and Ausubel et al., Current Protocols in Molecular Biology, Green Publishing Associates and Wiley
- 30 Interscience, N.Y. (2001). Alternatively, the nucleic acid molecules and vectors of the invention can be reconstituted into liposomes for delivery to target cells. The vectors containing the nucleic acid molecules of the invention can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for
- 35 prokaryotic cells, whereas, e.g., calcium phosphate or DEAE-Dextran mediated

5 transfection or electroporation may be used for other cellular hosts; see Sambrook, supra.

Such vectors may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions. Preferably, the nucleic acid molecule of the invention is operatively linked to
10 expression control sequences allowing expression in prokaryotic or eukaryotic cells. Expression of said polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and, optionally, a
15 poly-A signal ensuring termination of transcription and stabilization of the transcript, and/or an intron further enhancing expression of said polynucleotide. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the PL, lac,
20 trp or tac promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may
25 also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. In this context, suitable expression vectors are
30 known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3, the Echo™ Cloning System (Invitrogen), pSPORT1 (GIBCO BRL) or pRevTet-On/pRevTet-Off or pCI (Promega).

Preferably, the expression control sequences will be eukaryotic promoter systems in
35 vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used.

- 5 As mentioned above, the vector of the present invention may also be a gene transfer or targeting vector. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable vectors and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, *Nature Medicine* 2 (1996), 534-539; Schaper, *Circ. Res.* 79 (1996), 911-919; Anderson, *Science* 256 (1992), 808-813; Isner, *Lancet* 348 (1996), 370-374; Muhlhauser, *Circ. Res.* 77 (1995), 1077-1086; Wang, *Nature Medicine* 2 (1996), 714-716; WO94/29469; WO 97/00957, Schaper, *Current Opinion in Biotechnology* 7 (1996), 635-640, or Kay et al. (2001) *Nature Medicine*, 7, 33-40) and references cited therein. The polynucleotides and vectors of the invention may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell. Gene therapy is envisaged with the wild-type nucleic acid molecule only.
- 20 The invention also relates to a primer or primer pair, wherein the primer or primer pair hybridizes under stringent conditions to the nucleic acid molecule of the present invention comprising nucleotide positions 3966 and/or 5205 SEQ ID NO:1 or to the complementary strand thereof. In a preferred embodiment, said primer has an adenine or a guanine residue in the position corresponding to position 3966 of the USF1 sequence. In another preferred embodiment, said primer has a cytosine or a thymine residue in the position corresponding to position 5205 of the USF1 sequence. The primer may bind to the coding (+) strand or to the non-coding (-) strand of the DNA double strand.
- 30 Preferably, the primers of the invention have a length of at least 14 nucleotides such as 17, 20 or 21 nucleotides. The fact that in one embodiment the target sequence of the primer is located 3' to the SNP is to ensure that the primer is actually useful for sequence analysis, i.e. that the elongated primer sequence actually contains the SNP. When a PCR reaction is performed, for example, usually two primers are involved, wherein one primer binds 3' of the SNP on the + strand and the other primer binds 3' of the SNP on the - strand.
- 35

5 In one embodiment, the primer actually binds to the position of the SNP. As a consequence, when binding is performed under stringent conditions, such a primer is useful to distinguish between different polymorphic variants as binding only occurs if the sequences of the primer and the target have full complementarity. It is further preferred that the primers have a maximum length of 24 nucleotides.

10 However, in particular cases it may be preferable to use primers with a maximum length of 30 or 35 nucleotides. Hybridization or lack of hybridization of a primer under appropriate conditions to a genome sequence comprising either position 3966 or position 5205 coupled with an appropriate detection method such as an elongation reaction or an amplification reaction may be used to differentiate

15 between the polymorphic variants and then draw conclusions with regard to, e.g., the predisposition of the person under investigation hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism. The present invention envisages two types of primers/primer pairs. One type hybridizes to a sequence comprising the mutant, i.e. disease-associated sequence. In other terms. One

20 nucleotide of the primer pairs with the guanine residue in position 3966 (or the cytosine residue of the complementary strand) or with the thymine residue in position 5205 (or the adenine residue in the complementary strand). The other type of primer is exactly complementary to a sequence of wild-type. Since hybridization conditions would preferably be chosen to be stringent enough, contacting of e.g. a

25 primer exactly complementary to the mutant sequence with a wild-type allele would not result in efficient hybridization due to the mismatch formation. After washing, no signal would be detected due to the removal of the primer.

Additionally, the invention relates to a non-human host transformed with the vector of the invention as described herein above. The host may either carry the mutant or

30 the wild-type sequence. Upon breeding etc. the host may be heterozygous or homozygous for one or both SNPs.

The host of the invention may carry the vector of the invention either transiently or stably integrated into the genome. Methods for generating the non-human host of the invention are well known in the art. For example, conventional transfection

35 protocols described in Sambrook et al., loc. cit., may be employed to generate transformed bacteria (such as *E. coli*) or transformed yeasts. The non-human host

- 5 of the invention may be used, for example, to elucidate the onset of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism.

In a preferred embodiment of the invention the non-human host is a bacterium, a yeast cell, an insect cell, a fungal cell, a mammalian cell, a plant cell, a transgenic animal or a transgenic plant.

- 10 Whereas *E. coli* is a preferred bacterium, preferred yeast cells are *S. cerevisiae* or *Pichia pastoris* cells. Preferred fungal cells are *Aspergillus* cells and preferred insect cells include *Spodoptera frugiperda* cells. Preferred mammalian cells are CHO cells, colon carcinoma and hepatoma cell lines showing expression of the USF1 transcription factor. However, also cell lines with very low expression of USF1,
15 including HeLa cells and the like or fibroblasts, might be particularly useful for specific experiments.

- A method for the production of a transgenic non-human animal, for example transgenic mouse, comprises introduction of the aforementioned polynucleotide or targeting vector into a germ cell, an embryonic cell, stem cell or an egg or a cell
20 derived therefrom. The non-human animal can be used in accordance with a screening method of the invention described herein. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., *Gene Targeting, A Practical Approach* (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using, e.g.,
25 Southern blots with an appropriate complementary nucleic acid molecule; see supra. A general method for making transgenic non-human animals is described in the art, see for example WO 94/24274. For making transgenic non-human organisms (which include homologously targeted non-human animals), embryonal stem cells (ES cells) are preferred. Murine ES cells, such as AB-1 line grown on
30 mitotically inactive SNL76/7 cell feeder layers (McMahon and Bradley, *Cell* 62:1073-1085 (1990)) essentially as described (Robertson, E. J. (1987) in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*. E. J. Robertson, ed. (Oxford: IRL Press), p. 71-112) may be used for homologous gene targeting. Other suitable ES lines include, but are not limited to, the E14 line (Hooper et al., *Nature* 326:292-295
35 (1987)), the D3 line (Doetschman et al., *J. Embryol. Exp. Morph.* 87:27-45 (1985)),

5 the CCE line (Robertson et al., Nature 323:445-448 (1986)), the AK-7 line (Zhuang
et al., Cell 77:875-884 (1994)). The success of generating a mouse line from ES
cells bearing a specific targeted mutation depends on the pluripotency of the ES
cells (i. e., their ability, once injected into a host developing embryo, such as a
10 blastocyst or morula, to participate in embryogenesis and contribute to the germ
cells of the resulting animal). The blastocysts containing the injected ES cells are
allowed to develop in the uteri of pseudopregnant nonhuman females and are born
as chimeric mice. The resultant transgenic mice are chimeric for cells having the
desired nucleic acid molecule are backcrossed and screened for the presence of the
correctly targeted transgene (s) by PCR or Southern blot analysis on tail biopsy
15 DNA of offspring so as to identify transgenic mice heterozygous for the nucleic acid
molecule of the invention.

The transgenic non-human animals may, for example, be transgenic mice, rats,
hamsters, dogs, monkeys (apes), rabbits, pigs, or cows. Preferably, said transgenic
non-human animal is a mouse. The transgenic animals of the invention are, inter
20 alia, useful to study the phenotypic expression/outcome of the nucleic acids and
vectors of the present invention. Furthermore, the transgenic animals of the present
invention are useful to study the developmental expression of the USF1 gene and of
its role for onset of hyperlipidemia and/or dyslipidemia and/or defective
carbohydrate metabolism, for example in the rodent intestine. It is furthermore
25 envisaged, that the non-human transgenic animals of the invention can be
employed to test for therapeutic agents/compositions or other possible therapies
which are useful to hyperlipidemia and/or dyslipidemia and/or defective
carbohydrate metabolism.

The present invention also relates to a pharmaceutical composition comprising
30 USF1 or a fragment thereof, a nucleic acid molecule encoding USF1 or a fragment
thereof, or an antibody specific for USF1.

The components of the pharmaceutical composition of the invention may be
combined with a pharmaceutically acceptable carrier and/or diluent and/or excipient.
Preferably, USF1 refers to any USF1 being capable of alleviating the disease
35 symptoms. Generally, USF1 will be of wild-type. However, in particular cases it

5 might also be useful to administer mutated USF1 having one or more point mutations, insertions, deletions and the like and showing increased or decreased function or activity. Also encompassed by the present invention are chemically modified molecules which improve uptake or stability of a polypeptide.

10 Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Compositions comprising such carriers can be formulated by well known conventional methods. These pharmaceutical compositions can be administered to the subject at a suitable dose.

15 Administration of the suitable compositions may be effected by different ways, e.g., by intravenous, intraperitoneal, subcutaneous, intramuscular, topical, intradermal, intranasal or intrabronchial administration. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time

20 and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000 μg of nucleic acid for expression or for inhibition of expression; however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately 10^6 to 10^{12} copies of the DNA

25 molecule. Progress can be monitored by periodic assessment. The compositions of the invention may be administered locally or systemically. Administration will generally be parenterally, e.g., intravenously; DNA may also be administered directly to the target site, e.g., by biolistic delivery to an internal or external target

30 site or by catheter to a site in an artery. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or

35 suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated

5 Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

10 Additionally, the invention relates to a diagnostic composition comprising a nucleic acid molecule encoding USF1 or a fragment thereof, the nucleic acid molecule as described herein above, the vector as described herein above, the primer or primer pair as described herein above or an antibody specific for USF1.

15 The diagnostic composition is useful for assessing the genetic status of a person with respect to his or her predisposition to develop hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism or with regard to the diagnosis of the acute condition. The various possible components of the diagnostic composition may be packaged in one or more vials, in a solvent or otherwise such as in lyophilized form. If dissolved in a solvent, the diagnostic composition is preferably cooled to at least +8°C to +4°C. Freezing may be preferred in other
20 instances.

The present invention also relates to a method for testing for the presence or predisposition of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism, comprising analyzing a sample obtained from a prospective patient or from a person suspected of carrying such a predisposition for the presence of a
25 wild-type or variant allele of the USF1 gene. Preferably, said variant comprises an SNP at position 3966 and/or at position 5205 of the USF1 gene in a homozygous or heterozygous state. In varying embodiments, it may be tested either for the presence of the wild-type sequence(s) or of the mutant sequence(s). It is in accordance with the present invention that a guanine residue in position 3966 of the
30 USF1 gene indicates the presence of a disease-associated allele, whereas an adenine residue in the same position of the USF1 gene is indicative for the healthy allele. Likewise, a cytosine residue in position 5205 of the USF1 gene indicates the presence of a disease-associated allele, whereas a thymine residue is indicative for the healthy allele.

5 The method of the invention is useful for detecting the genetic set-up of said person/patient and drawing appropriate conclusions whether a condition from which said patient suffers is hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism. Alternatively, it may be assessed whether a person not suffering from a condition carries a predisposition to hyperlipidemia and/or
10 dyslipidemia and/or defective carbohydrate metabolism. With regard to position 5205 in exon 11 of the USF1 gene, only if cytosine is found in a homozygous or heterozygous state, a condition would be diagnosed as hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism or a corresponding predisposition would be manifest. On the other hand, if thymine is found in a
15 homozygous state, then it may be concluded that a condition from which a patient suffers is not related to hyperlipidemia or dyslipidemia and/or defective carbohydrate metabolism and further, that the patient does not carry a predisposition to develop this condition. The situation is similar and essentially the same conclusions apply for the analysis of the SNP in position 3966: With regard to position 3966 in intron 7 of
20 the USF1 gene, only if guanine is found in a homozygous or heterozygous state, a condition would be diagnosed as hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism or a corresponding predisposition would be manifest. On the other hand, if an adenine is found in a homozygous state, then it may be concluded that a condition from which a patient suffers is not related to
25 hyperlipidemia or dyslipidemia and/or defective carbohydrate metabolism and further, that the patient does not carry a predisposition to develop this condition.

In a preferred embodiment of the method of the invention said testing comprises hybridizing the complementary nucleic acid molecule as described herein above which is complementary to the nucleic acid molecule contributing to or indicative of
30 hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism or the nucleic acid molecule as described herein above which is complementary to the wild-type sequence as a probe under (highly) stringent conditions to nucleic acid molecules comprised in said sample and detecting said hybridization, wherein said complementary nucleic acid molecule comprises the sequence position containing
35 the SNP.

5 Again, depending on the nucleic acid probe used, either wild-type or mutant sequences (i.e. sequences contributing to or indicative of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism) would be detected. It is understood that hybridization conditions would be chosen such that a nucleic acid molecule complementary to wild-type sequences would not or essentially not
10 hybridize to the mutant sequence. Similarly, a nucleic acid molecule complementary to the mutant sequence would not or would not essentially not hybridize to the wild-type sequence. In order to differentiate between results obtained from homozygous and heterozygous genotypes in the hybridization methods of the invention, one can for example monitor/detect the strength/intensity of the respective detection signal
15 after the hybridization. To differentiate between wild-type homozygous, heterozygous and/or mutant homozygous alleles in the hybridization methods of the invention, internal control samples of the corresponding genotypes will be included in the analysis.

20 In a further preferred embodiment, the method of the invention further comprises digesting the product of said hybridization with a restriction endonuclease or subjecting the product of said hybridization to digestion with a restriction endonuclease and analyzing the product of said digestion.

25 This preferred embodiment of the invention allows by convenient means, the differentiation between an effective hybridization and a non-effective hybridization. For example, if the DNA sequence adjacent to position 3966 or position 5205 comprises an endonuclease restriction site, the hybridized product will be cleavable by an appropriate restriction enzyme upon an effective hybridization whereas a lack of hybridization will yield no double-stranded product or will not comprise the recognizable restriction site and, accordingly, will not be cleaved. Suitable restriction
30 enzymes may be found, for example, by the use of the program Webcutter. The analysis of the digestion product can be effected by conventional means, such as by gel electrophoresis which may be optionally combined by the staining of the nucleic acid with, for example, ethidium bromide. Combinations with further techniques such as Southern blotting are also envisaged.

5 Detection of said hybridization may be effected, for example, by an anti-DNA double-strand antibody or by employing a labeled oligonucleotide. Conveniently, the method of the invention is employed together with blotting techniques such as Southern or Northern blotting and related techniques. Labeling may be effected, for example, by standard protocols and includes labeling with radioactive markers,
10 fluorescent, phosphorescent, chemiluminescent, enzymatic labels, etc. The label can be located at the 5' and/or 3' end of the nucleic acid molecule or be located at an internal position. Preferred labels include, but are not limited to, fluorochromes, e.g. Carboxyfluorescein (FAM) and 6-carboxy-X-rhodamine (ROX), fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-
15 carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, e.g. ^{32}P , ^{35}S , ^3H ; etc. The label may also be a two stage system, where the probe is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin,
20 specific antibodies, etc., where the binding partner is conjugated to a detectable label.

In accordance with the above, in another preferred embodiment of the method of the invention said probe is detectably labeled, e.g. by the methods and with the labels described herein above.

25 In yet another preferred embodiment of the method of the invention said testing comprises determining the nucleic acid sequence of at least a portion of the nucleic acid molecule as described herein above, said portion comprising the position of the SNP. Determination of the nucleic acid molecule may be effected in accordance with one of the conventional protocols such as the Sanger or Maxam/Gilbert
30 protocols (see Sambrook et al., loc. cit., for further guidance).

In a further preferred embodiment of the method of the invention the determination of the nucleic acid sequence is effected by solid-phase minisequencing. Solid-phase minisequencing is based on quantitative analysis of the wild type and mutant nucleotide in a solution. First, the genomic region containing the mutation is
35 amplified by PCR with one biotinylated and non-biotinylated primer where the

5 biotinylated primer is attached to a streptavidin (SA) coated plate. The PCR-product is denatured to a single stranded form to allow a minisequencing primer to bind to this strand just before the site of the mutation. The tritium (H3) or fluorescence labeled mutated and wild type nucleotides together with nonlabeled dNTPs are added to the minisequencing reaction and sequenced using Taq-polymerase. The
10 result is based on the amount of wild type and mutant nucleotides in the reaction measured by beta counter or fluorometer and expressed as an R-ratio. See also Syvänen AC, Sajantila A, Lukka M. Am J Hum Genet 1993: 52,46-59 and Suomalainen A and Syvanen AC. Methods Mol Biol 1996;65:73-79.

15 A preferred embodiment of the method of the invention further comprises, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid molecule. Preferably, amplification is effected by polymerase chain reaction (PCR). Other amplification methods such as ligase chain reaction may also be employed.

20 In a preferred embodiment of the method of the invention said testing comprises carrying out an amplification reaction wherein at least one of the primers employed in said amplification reaction is the primer as described herein above or belongs to the primer pair as described herein above, comprising assaying for an amplification product. In this embodiment and depending on the information the investigator/physician wishes to obtain, primers hybridizing either to the wild-type or
25 mutant sequences may be employed. In a particularly preferred embodiment, at least one of the primers will actually bind to the position of the SNP. As a consequence, when binding is performed under stringent conditions, such a primer is useful to distinguish between different polymorphic variants as binding only occurs if the sequences of the primer and the target have full complementarity.

30 The method of the invention will result in an amplification of only the target sequence, if said target sequence carries a sequence exactly complementary to the primer used for hybridization. This is because the oligonucleotide primer will under preferably (highly) stringent hybridization conditions not hybridize to the wild-type/mutant sequence – depending which type of primer is used – (with the
35 consequence that no amplification product is obtained) but only to the exactly

- 5 matching sequence. Naturally, combinations of primer pairs hybridizing to both SNPs may be used. In this case, the analysis of the amplification products expected (which may be no, one, two, three or four amplification product(s) if the second, non-differentiating primer is the same for each locus) will provide information on the genetic status of both positions 3966 and 5205.
- 10 In a preferred embodiment of the method of the invention said amplification is effected by or said amplification is the polymerase chain reaction (PCR). The PCR is well established in the art. Typical conditions to be used in accordance with the present invention include for example a total of 35 cycles in a total of 50µl volume exemplified with a denaturation step at 93° C for 3 minutes; an annealing step at 55°
- 15 C for 30 seconds; an extension step at 72° C for 75 seconds and a final extension step at 72° C for 10 minutes.

The present invention further relates to a method for testing for the presence or predisposition of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism comprising assaying a sample obtained from a human for the amount of

20 USF1 contained in said sample. The amount of USF1 can be determined by any suitable method. Preferably, the amount of USF1 is determined by contacting the sample, i.e. USF1 contained in the sample, with an antibody or aptamer or a derivative thereof, which is specific for USF1. For example, the sample containing USF1 may be analyzed in a Western blot or in a RIA assay. In this context a weaker

25 staining for the presence of the antigen of the invention compared to homozygous wild type control samples (comprising two persistent alleles) is indicative for the heterozygous wild type (one persistent allele and one disease-associated allele), whereas for the homozygous disease state no staining or a reduced staining is expected if the appropriate antibody is used. Preferably, the method of the invention

30 is performed in the presence of control samples corresponding to all three possible allelic combinations as internal controls. Testing may be carried out with an antibody or aptamer etc. specific for the wild-type or specific for the mutant sequence. Testing for binding may, again, involve the employment of standard techniques such as ELISAs; see, for example, Harlow and Lane⁵³, loc. cit. The term "antibody" as

35 used throughout the invention refers to monoclonal antibodies, polyclonal antibodies, single chain antibodies, or a fragment thereof. Preferably the antibody is

5 specific for USF1 or for wild-type or disease-associated USF1. The antibodies may be bispecific antibodies, humanized antibodies, synthetic antibodies, antibody fragments, such as Fab, a F(ab₂)', Fv or scFv fragments etc., or a chemically modified derivative of any of these (all comprised by the term "antibody"). Monoclonal antibodies can be prepared, for example, by the techniques as originally
10 described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, Meth. Enzymol. 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals with modifications developed by the art. Antibodies may be labelled by using any of the labels described in the present invention.

15 In a preferred embodiment of the method of the invention said antibody or aptamer is detectably labeled. Whereas the aptamers are preferably radioactively labeled with ³H or ³²P or with a fluorescent marker, the antibody may either be labeled in a corresponding manner (with ¹³¹I as the preferred radioactive label) or be labeled with a tag such as His-tag, FLAG-tag or myc-tag.

20 In a further preferred embodiment of the method of the invention the test is an immuno-assay.

In another preferred embodiment of the method of the invention said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human
25 tissue.

In an additional preferred embodiment of the method of the invention said nucleic acid molecule from said sample is fixed to a solid support.

Fixation of the nucleic acid molecule to a solid support will allow an easy handling of the test assay and furthermore, at least some solid supports such as chips, silica
30 wafers or microtiter plates allow for the simultaneous analysis of larger numbers of samples. Ideally, the solid support allows for an automated testing employing, for example, robotics devices.

- 5 In a particularly preferred embodiment of the method of the invention said solid support is a chip, a silica wafer, a bead or a microtiter plate.

The methods of the present invention may be performed *ex vivo*, *in vitro* or *in vivo*.

- 10 The present invention also relates to the use of a nucleic acid molecule encoding USF1, the nucleic acid molecule as described herein above, or of USF1 polypeptide for the analysis of the presence or predisposition of hyperlipidemia, dyslipidemia and/or defective carbohydrate metabolism. The nucleic acid molecule simultaneously allows for the analysis of the absence of the condition or the predisposition to the condition, as has been described in detail herein above. In particular cases, it may be possible to use USF1 polypeptides for testing. This may
15 be, for example, in cases when expression of USF1 results in an autoimmune response against USF1. In such cases it will be possible, by using USF1 polypeptides, to monitor patients by detecting antibodies directed against USF1. Such assays can, for example, be based on the western blotting technique or by performing (radio)immunoprecipitations.

- 20 In addition, the present invention relates to the use of USF1 or a fragment thereof, a nucleic acid molecule encoding USF1 and/or comprising at least the wild-type sequence of intron 7 and/or exon 11 of USF1, for the preparation of a pharmaceutical composition for the treatment of hyperlipidemias and/or dyslipidemias, including familial combined hyperlipidemia (FCHL),
25 hypercholesterolemia, hypertriglyceridemia, hypoalphalipoproteinemia, hyperapobetalipoproteinemia (hyperapoB) and/or familial dyslipidemic hypertension (FDH), coronary heart disease, type II diabetes, atherosclerosis or metabolic syndrome. Any of the diseases mentioned in the present invention can be treated by administering to a patient USF1 in an amount and quality sufficient to ameliorate the
30 symptoms of the disease. If for example the disease symptoms are created by a reduced amount of USF1 in the patient, administration of USF1 to the patient will compensate for the reduced USF1 of the patient. USF1 may be provided to the patient as such, i.e. as the polypeptide. Alternatively, a nucleic acid molecule encoding USF1 can be administered. Preferably, USF1 is a full length wild-type
35 polyprotein. However, in particular cases it might also be useful to administer

5 mutated USF1 having one or more point mutations, insertions, deletions and the like
and showing increased or decreased function or activity. Also encompassed by the
present invention are chemically modified molecules which improve uptake or
stability of a polypeptide. Gene therapy approaches have been discussed herein
above in connection with the vector of the invention and equally apply here. It is of
10 note that in accordance with this invention, also fragments of the nucleic acid
molecules as defined herein above may be employed in gene therapy approaches.
Said fragments comprise the nucleotide at position 3966 as or position 5205 of the
USF1 gene. Preferably, said fragments comprise at least 200, at least 250, at least
300, at least 400 and most preferably at least 500 nucleotides. In a preferred
15 embodiment of the use of the invention said gene therapy treats or prevents
hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism.

Finally, the present invention relates to a kit comprising the nucleic acid molecule,
the primer or primer pair and/or the vector of the present invention in one or more
containers.

References

1. Goldstein, J.L., Schrott, H.G., Hazzard, W.R., Bierman, E.L. & Motulsky, A.G. Hyperlipidemia in coronary heart disease II. Genetic analysis of lipid levels in 176 families and delineation of a new inherited disorder, combined hyperlipidemia. *J. Clin. Invest.* **52**, 1544–1568 (1973).
2. Nikkilä, E.A. & Aro, A. Family study of serum lipids and lipoproteins in coronary heart disease. *Lancet* **1**, 954–959 (1973).
3. Genest, J.J. Jr. *et al.* Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* **85**, 2025–2033 (1992).
4. Pajukanta, P. *et al.* Linkage of familial combined hyperlipidemia to chromosome 1q21–q23. *Nat. Genet.* **18**, 369–373 (1998).
5. Coon, H. *et al.* Replication of linkage of familial combined hyperlipidemia to chromosome 1q with additional heterogeneous effect of apolipoprotein A-I/C-III/A-IV locus: the NHLBI family heart study. *Arterioscler. Thromb. Vasc. Biol.* **20**, 2275–2280 (2000).
6. Pei, W. *et al.* Support for linkage of familial combined hyperlipidemia to chromosome 1q21–q23 in Chinese and German families. *Clin. Genet.* **57**, 29–34 (2000).
7. Allayee, A. *et al.* Locus for Elevated Apolipoprotein B Levels on Chromosome 1p31 in Families with Familial Combined Hyperlipidemia. *Circ. Res.* **90**, 926–931 (2002).
8. Aouizerat, B.E. *et al.* A genome scan for familial combined hyperlipidemia reveals evidence of linkage with a locus on chromosome 11. *Am. J. Hum. Genet.* **65**, 397–412 (1999).
9. Pajukanta, P. *et al.* Genomewide scan for familial combined hyperlipidemia genes in Finnish families, suggesting multiple susceptibility loci influencing triglyceride, cholesterol and apolipoprotein B levels. *Am. J. Hum. Genet.* **64**, 1453–1463 (1999).
10. Elbein, S.C., Hoffman, M.D., Teng, K., Leppert, M.F. & Hasstedt, S.J. A genome-wide search for type 2 diabetes susceptibility genes in Utah Caucasians. *Diabetes* **48**, 1175–1182 (1999).
11. Hanson, R.L. *et al.* An autosomal genomic scan for loci linked to type II diabetes mellitus and body-mass index in Pima Indians. *Am. J. Hum. Genet.* **63**, 1130–1138 (1998).
12. Vionnet, N., Hani, El.-H., Dupont, S., Gallina, S., Francke, S. & Dotte, S. Genomewide search for type 2 diabetes-susceptibility genes in French whites: evidence for a novel susceptibility locus for early-onset diabetes on

- 5 chromosome 3q-qter and independent replication of a type 2-diabetes locus on chromosome 1q21-q24. *Am. J. Hum. Genet.* **67**, 1470-1480 (2000).
13. Wiltshire, S. et al. A genomewide scan for loci predisposing to type 2 diabetes in a U.K. population (the Diabetes UK Warren 2 Repository): analysis of 573 pedigrees provides independent replication of a susceptibility locus on chromosome 1q. *Am. J. Hum. Genet.* **69**, 553-569 (2001).
- 10 14. Hsueh, W.C. et al. Genome-wide and fine-mapping linkage studies of type 2 diabetes and glucose traits in the Old Order Amish: evidence for a new diabetes locus on chromosome 14q11 and confirmation of a locus on chromosome 1q21-q24. *Diabetes* **52**, 550-507 (2003).
- 15 15. Watanabe, R.M. et al. The Finland-United States investigation of non-insulin-dependent diabetes mellitus genetics (FUSION) study. II. An autosomal genome scan for diabetes-related quantitative-trait loci. *Am. J. Hum. Genet.* **67**, 1186-1200 (2000).
- 20 16. Castellani, L.W. et al. Mapping a gene for combined hyperlipidaemia in a mutant mouse strain. *Nat. Genet.* **18**, 374-377 (1998).
17. Bodnar, J.S. et al. Positional cloning of the combined hyperlipidemia gene Hyplip1. *Nat. Genet.* **30**, 110-116 (2002).
- 25 18. Salero, E., Gimenez, C. & Zafra, F. Identification of a non-canonical E-box motif as a regulatory element in the proximal promoter region of the apolipoprotein E gene. *Biochem. J.* **370**, 979-986 (2003).
19. Portois, L., Tastenoy, M., Viollet, B. & Svoboda, M. Functional analysis of the glucose response element of the rat glucagon receptor gene in insulin-producing INS-1 cells. *Biochim. Biophys. Acta.* **1574**, 175-186 (2002).
- 30 20. Yang, X.P. et al. The E-box motif in the proximal ABCA1 promoter mediates transcriptional repression of the ABCA1 gene. *J. Lipid. Res.* **43**, 297-306 (2002).
21. Smih, F. et al. Transcriptional regulation of adipocyte hormone-sensitive lipase by glucose. *Diabetes* **51**, 293-300 (2002).
- 35 22. Casado, M., Vallet, V.S., Kahn, A. & Vaulont, S. Essential role in vivo of upstream stimulatory factors for a normal dietary response of the fatty acid synthase gene in the liver. *J. Biol. Chem.* **274**, 2009-2013 (1999).
23. Ribeiro, A., Pastier, D., Kardassis, D., Chambaz, J. & Cardot, P. Cooperative binding of upstream stimulatory factor and hepatic nuclear factor 4 drives the transcription of the human apolipoprotein A-II gene. *J. Biol. Chem.* **274**, 1216-1225 (1999).
- 40 24. Iynedjian, P.B. Identification of upstream stimulatory factor as transcriptional activator of the liver promoter of the glucokinase gene. *Biochem. J.* **333**, 705-712 (1998).

- 5 25. Soro, A., Jauhiainen, M., Ehnholm, C. & Taskinen, M.R. Determinants of low HDL levels in familial combined hyperlipidemia. *J. Lipid Res.* **44**, 1536-1544 (2003).
26. Risch, N. & Teng, J. The relative power of family-based and case-control designs for linkage disequilibrium studies of complex human diseases I. DNA pooling. *Genome Res.* **8**, 1273-1288 (1998).
- 10 27. Terwilliger, J.D. & Ott, J. A haplotype-based 'haplotype relative risk' approach to detecting allelic associations. *Hum. Hered.* **42**, 337-346 (1992).
28. Spielman, R.S., McGinnis, R.E. & Ewens, W.J. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am. J. Hum. Genet.* **52**, 506-516 (1993).
- 15 29. Sinsheimer, J.S., Blangero, J. & Lange, K. Gamete competition models. *Am. J. Hum. Genet.* **66**, 1168-1172 (2000).
30. Peltonen, L., Pekkarinen, P. & Aaltonen, J. Messages from an isolate: lessons from the Finnish gene pool. *Biol. Chem.* **376**, 697-704 (1995).
- 20 31. Rioux, J.D. *et al.* Genetic variation in the 5q31 cytokine gene cluster confers susceptibility to Crohn disease. *Nat. Genet.* **29**, 223-228 (2001).
32. Vallet, V.S. *et al.* Differential roles of upstream stimulatory factors 1 and 2 in the transcriptional response of liver genes to glucose. *J. Biol. Chem.* **273**, 20175-20179 (1998).
- 25 33. Wang, D. & Sul, H.S. Upstream stimulatory factor binding to the E-box at -65 is required for insulin regulation of the fatty acid synthase promoter. *J. Biol. Chem.* **272**, 26367-26374 (1997).
- 30 34. Pan, L. *et al.* Critical roles of a cyclic AMP responsive element and an E-box in regulation of mouse renin gene expression. *J. Biol. Chem.* **276**, 45530-45538 (2001).
- 35 35. Yanai, K., Saito, T., Hirota, K., Kobayashi, H., Murakami, K. & Fukamizu, A. Molecular variation of the human angiotensinogen core promoter element located between the TATA box and transcription initiation site affects its transcriptional activity. *J. Biol. Chem.* **272**, 30558-30562 (1997).
36. Barton, E.S. *et al.* Junction adhesion molecule is a receptor for reovirus. *Cell* **104**, 441-451 (2001).
37. Ostermann, G., Weber, K.S., Zernecke, A., Schroder, A. & Weber, C. JAM-1 is a ligand of the beta(2) integrin LFA-1 involved in transendothelial migration of leukocytes. *Nat. Immunol.* **3**, 151-158 (2002).
- 40 38. Enattah, N.S., Sahi, T., Savilahti, E., Terwilliger, J.D., Peltonen, L. & Jarvela, I. Identification of a variant associated with adult-type hypolactasia. *Nat. Genet.* **30**, 233-237 (2002).

- 5 39. Vakkilainen, J., Jauhiainen, M., Ylitalo, K., Nuotio, I.O., Viikari, J.S., Ehnholm, C. & Taskinen, M.R. LDL particle size in familial combined hyperlipidemia: effects of serum lipids, lipoprotein-modifying enzymes, and lipid transfer proteins. *J. Lipid Res.* **43**, 598-603 (2002).
- 10 40. Pielberg, G., Olsson, C., Syvänen, A.-C. & Andersson, L. Unexpectedly High Allelic Diversity at the *KIT* Locus Causing Dominant White Color in the Domestic Pig. *Genetics* **160**, 305-311 (2002).
41. The Gene Ontology Consortium Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25-29 (2000).
- 15 42. Hosack, D.A., Dennis, G. Jr., Sherman, B.T., Lane, H.C., Lempicki, R.A. Identifying biological themes within lists of genes with EASE. *Genome Biol.* **4**, R70.1-R70.8 (2003).
43. Lathrop, G.M., Lalouel, J.-M., Julier, C.A. & Ott, J. Strategies for multilocus linkage analysis in humans. *Proc. Natl. Acad. Sci. USA* **81**, 3443-3446 (1984).
- 20 44. Cottingham, R.W., Jr, Idury, R.M. & Schäffer, A.A. Faster sequential genetic linkage computations. *Am. J. Hum. Genet.* **53**, 252-263 (1993).
45. Schaffer, A.A., Gupta, S.K., Shriram, K. & Cottingham, R.W., Jr. Avoiding recomputation in linkage analysis. *Hum. Hered.* **44**, 225-237 (1994).
- 25 46. Göring, H.H. & Terwilliger, J.D. Gene mapping in the 20th and 21st centuries: statistical methods, data analysis, and experimental design. *Hum. Biol.* **72**, 63-132 (2000a).
47. Göring, H.H. & Terwilliger, J.D. Linkage analysis in the presence of errors III: marker loci and their map as nuisance parameters. *Am. J. Hum. Genet.* **66**, 1298-1309 (2000b).
- 30 48. Terwilliger, J.D. A powerful likelihood method for the analysis of linkage disequilibrium between trait loci and one or more polymorphic marker loci. *Am. J. Hum. Genet.* **56**, 777-787 (1995).
49. Laird, N., Horvath, S. & Xu, X. Implementing a unified approach to family based tests of association. *Genet. Epidemiol.* **19**, 36-42 (2000).
- 35 50. Martin, E.R., Bass, M.P., Gilbert, J.R., Pericak-Vance, M.A., & Hauser ER. *Genet. Epidemiol.* (2003, in press)

5 The figures show:

Figure 1: Schematic overview of the associated region on 1q21. Genes for which we genotyped SNPs as well as the locations of the peak linkage markers *D1S104* and *D1S1677* (Pajukanta et al. 1998) are shown in the uppermost part. The genes indicated in bold were also sequenced. Next part shows the SNPs genotyped for *JAM1* and *USF1* (see Table 2 for distances, rs numbers and LD clusters of these SNPs). The second to lowest part indicates the SNPs associated with TGs in men, and the lowest part the SNPs associated with FCHL and TGs in all family members.

Figure 2: Distribution of genes according to functional category for the 16 up-regulated and 60 down-regulated genes for which annotation information for the gene ontology (GO) class Biological process was available. Only categories scoring a statistically significant EASE-score (<0.05) for over-representation are shown. Complete results of the EASE analysis including the corresponding EASE scores (p-values) and the lists of genes in every significant category are given in the Supplementary Table 3a-b.

Figure 3a: Intron 7 of *USF1* harbors the 60-bp sequence shared by the 91 *USF1*-similarity genes. Parts (2-61 bp and 137-196 bp) of the AluSx repeat in intron 7 of *USF1* have sequence similarities with the mouse B1 repeat. A total of 91 human genes, including *USF1*, have this 60-bp part of AluSx located either on the coding strand (43 genes) or on the opposite strand (48 genes). These 91 genes are listed in the Supplementary Table 4.

Figure 3b: Transcription efficiency of a 268-bp region in intron 7 of *USF1* containing the critical 60-bp sequence and the *usf1s2* SNP (see Fig. 3a). DNAs from one homozygous susceptibility carrier (haplotype 1-1)

5 and one homozygous non-carrier (2-2) were cloned to the SEAP
reporter system in both forward and reverse orientations. HC for and
HC rev indicate constructs of a haplotype carrier (1-1) DNA in forward
and reverse orientations; HNC for and HNC rev indicate constructs of
10 a haplotype non-carrier (2-2) DNA in forward and reverse orientations.
Culture media from cells transfected with the pSEAP2-Basic vector
was used as a negative control (Neg) and culture media from cells
transfected with the pSEAP2-Control vector as a positive control (Pos),
respectively. The monitoring of the SEAP protein was performed 48
and 72 hours post-transfection. Error bars represent SD of one
15 experiment done in triplicate. The size of the bar indicates the increase
in transcriptional activity when compared to the negative control which
is set to 1.

- 5 The examples illustrate the invention.

EXAMPLE 1: EXPERIMENTAL OUTLINE

All analyzed FCHL families had a proband with severe CHD and lipid phenotype, and on average 5-6 FCHL affected family members. These FCHL families exhibiting extreme and well-defined disease phenotypes were analyzed to identify the underlying gene contributing to FCHL on 1q21. We selected a regional candidate gene approach and sequenced four functionally relevant regional candidate genes on 1q21. The *TXNIP*, *USF1*, retinoid X receptor gamma (*RGRG*), and apolipoprotein A2 (*APOA2*) genes were sequenced to identify all possible variants. Of these, *TXNIP* initially represented the most promising positional candidate gene, because it has been shown to underlie the combined hyperlipidemia phenotype in mice¹⁷. The three additional regional genes were selected for sequencing based on their functional candidacy and close location (< 2.5 Mb) to the original peak linkage markers, *D1S104* and *D1S1677* (Fig.1). In parallel, we employed a functionally unbiased, genetic approach, where an initial set of SNPs for genes around the peak linkage markers were tested for association. A total of 60 SNPs were genotyped for 26 genes on 1q21. Fifty of these SNPs were located within 5.8 Mb, flanking *D1S104* and *D1S1677*. All 60 SNPs were genotyped in 238 family members of 42 FCHL families, including the 31 families of the original linkage study⁴, and 10 most promising SNPs in the extended sample of 721 family members from 60 FCHL families (see below). The results of the 60 SNPs are shown in the Supplementary Table 1.

5 SUPPLEMENTARY TABLE 1.

RESULTS OF THE 60 GENOTYPED SNPS OBTAINED IN TWO-POINT LINKAGE AND ASSOCIATION ANALYSES. A TOTAL OF FOUR TRAITS WERE ANALYZED: ALL INDIVIDUALS FOR THE FCHL AND TG TRAITS, AS WELL AS AFFECTED MALES FOR THE FCHL AND TG TRAITS.

Lod scores were obtained in two-point linkage analysis (see methods for details) and p-values in the association analysis using the HHRR test. Nf indicates not found in dbSNP or Celera databases. The SNP information for these SNPs will be submitted to the public database (dbSNP). SNPs indicated in bold were genotyped in the 60 extended FCHL families. All other results were obtained in the 42 nuclear FCHL families. P-values less than 0.05 are also shown in bold, whereas ns indicates non-significant (p-value greater than 0.05).

10

Gene	SNP	Distance in bp	FCHL		FCHL men		TG		TG men	
			Linkage	HHRR	Linkage	HHRR	Linkage	HHRR	Linkage	HHRR
			Lod	P-value	Lod	P-value	Lod	P-value	Lod	P-value
TXNIP	rs2236567	425	0.4	ns	0.0	ns	0.2	ns	0.3	ns
	Nf	1272	0.3	ns	0.0	ns	0.2	ns	0.0	ns
TXNIP	rs9245	3039	0.3	ns	0.1	ns	0.4	ns	0.8	ns
TXNIP	rs7211	8869064	0.6	ns	0.1	ns	0.3	ns	0.0	ns
MUC1	rs1611774	4214	0.2	ns	0.1	ns	1.1	ns	1.4	ns
MUC1	rs4072037	22637	0.0	ns	0.0	ns	0.0	ns	0.1	ns

GBA	rs1800473	1661529	0.4	ns	0.0	ns	0.5	ns	0.2	ns
NTRK1	rs6334	2762	0.2	ns	0.0	ns	0.2	ns	0.1	ns
NTRK1	rs6337	2326359	0.0	ns	0.0	ns	0.2	ns	0.1	ns
FY	rs12075	507737	1.0	ns	1.0	ns	1.5	ns	0.8	ns
CRP	rs1130864	367299	0.0	ns	0.0	ns	0.2	ns	0.3	ns
KCNJ9	rs4656876	521	0.0	ns	0.0	ns	0.4	ns	0.1	ns
KCNJ9	rs2180752	7051	0.0	ns	0.0	ns	0.0	ns	0.0	ns
KCNJ9	rs2737705	288	0.0	ns	0.0	ns	0.4	ns	0.6	ns
KCNJ9	rs2753268	302	0.0	ns	0.0	ns	0.2	ns	1.0	ns
KCNJ9	Nf	38761	0.1	ns	0.0	ns	0.2	ns	0.5	ns
ATP1A2	rs2295623	12474	0.0	ns	0.0	ns	0.0	ns	0.0	ns
ATP1A2	Nf	71117	0.0	ns	0.0	ns	0.0	ns	0.0	ns
PEA15	rs680083	66279	0.0	ns	0.0	ns	0.0	ns	0.0	ns
PXF	rs10594	56231	0.4	ns	0.1	ns	0.2	ns	0.4	ns
COPA	rs1802778	276599	0.0	ns	0.0	ns	0.0	ns	0.0	ns
SLAMF1	rs1061217	337887	0.5	ns	0.0	ns	0.2	ns	0.3	ns
ITLN2	rs1556519	24927	0.9	ns	0.1	ns	1.0	ns	1.1	ns

Flanking	rs2246485	25395	1.1	ns	0.0	ns	1.1	ns	0.3	ns
F11R										
F11R/f11rs1	rs836	1361	1.7	ns	0.1	ns	2.8	ns	0.9	0.03
F11R/f11rs2	rs790056	1561	0.9	ns	0.0	ns	0.5	ns	1.1	ns
F11R/f11rs3	rs790055	25608	0.7	ns	0.0	ns	0.4	ns	0.3	ns
F11R/f11rs4	hCV1459766	10572	1.8	ns	0.1	ns	2.7	ns	0.4	ns
F11R/f11rs5	rs4339888	1246	2.2	ns	0.1	ns	3.6	ns	0.6	0.02
F11R/f11rs6	rs3766383	951	0.0	ns	0.0	ns	0.0	ns	0.0	ns
USF1/usf1s1	rs3737787	1239	3.3	ns	0.3	0.04	2.1	ns	2.0	0.0009
USF1/usf1s2	rs2073658	12	2.0	ns	0.0	0.04	1.5	ns	1.8	0.002
USF1/usf1s3	rs2516841	17	1.3	ns	0.0	ns	1.8	ns	0.4	ns
USF1/usf1s4	rs2073657	526	0.4	ns	0.1	ns	1.1	ns	0.4	ns
USF1/usf1s5	rs2516840	1443	0.7	ns	0.0	ns	0.8	ns	0.2	ns
USF1/usf1s6	rs2073653	361	0.0	ns	0.0	ns	0.0	ns	0.0	ns
USF1/usf1s7	rs2516839	1249	0.7	ns	0.0	ns	2.1	ns	1.2	ns
USF1/usf1s8	rs2516838	279	0.1	ns	0.0	ns	0.4	ns	0.1	0.01
USF1/usf1s9	rs1556259	4391	0.0	ns	0.0	ns	0.0	ns	0.0	ns

LOC257106	rs3813609	5724	0.1	ns	0.0	ns	0.8	ns	0.1	ns
LOC257106	Nf	26087	0.1	ns	0.1	ns	0.1	ns	0.3	ns
LNIR	rs1467742	283	0.0	ns	0.0	ns	0.0	ns	0.0	ns
LNIR	rs1556257	2639	0.1	ns	0.1	ns	0.0	ns	0.0	ns
LNIR	rs4529727	87659	0.0	ns	0.0	ns	0.6	ns	0.2	ns
B4GALT3	rs6779	47461	0.1	ns	0.0	ns	0.3	ns	0.4	ns
FCER1G	rs3557	43	0.1	ns	0.0	ns	0.0	ns	0.0	ns
FCER1G	rs11421	2593	0.1	ns	0.0	ns	0.3	ns	0.0	ns
APOA2	Nf	34	0.3	ns	0.0	ns	0.6	ns	0.0	ns
APOA2	Nf	948	1.1	ns	0.0	ns	1.5	ns	0.0	ns
APOA2	rs5085	1172	0.1	ns	0.0	ns	0.0	ns	0.0	ns
APOA2	rs5082	645533	0.3	ns	0.1	ns	3.1	ns	2.1	ns
ATF6	CV67448	1196247	0.0	ns	0.0	ns	0.0	ns	0.0	ns
RGS5	rs15049	1412242	0.0	ns	0.1	ns	0.0	ns	0.0	ns
PBX1	rs2275558	122535	0.0	ns	0.0	ns	0.0	ns	0.0	ns
PBX1	rs1057756	164453	0.0	ns	0.2	ns	0.2	ns	0.1	ns
PBX1	rs14832	561444	0.1	ns	0.0	ns	1.3	ns	0.1	ns

RXRG	rs2134095	11733	0.3	ns	0.0	ns	1.2	ns	0.2	ns
RXRG	rs157870	242385	0.0	ns	0.0	ns	0.0	ns	0.0	ns
ALDH9A1	rs12670	307375	0.8	ns	0.0	ns	0.9	ns	0.0	ns
LMX1A	hCV3194556		0.9	ns	0.0	ns	0.9	ns	0.1	ns

5

EXAMPLE 2: *USF1* GENE AS A CANDIDATE GENE

We identified a total of 23 SNPs for the 5687 bp sequence of the *USF1* gene (Supplementary Table 2): Three of these were silent variants in exons, and the rest were located in the non-coding regions and in the putative promoter. Eight of the 23 SNPs were novel. Initially, we genotyped three SNPs for the *USF1* gene: *usf1s1* (exon 11), *usf1s2* (intron 7), and *usf1s7* (exon 2) (the corresponding rs numbers for the genotyped SNPs are given in Tables 2-3).

TABLE 1. MULTIPOINT HHRR AND GAMETE COMPETITION ANALYSES FOR THE SNPs *USF1s1* (=RS3737787) AND *USF1s2* (=RS2073658).

All values represent p-values for simultaneous analysis of both SNPs. *ns* indicates non-significant. The first presented p-values were obtained in 60 extended FCHL families and the values given in parentheses in 42 nuclear FCHL families. Gene dropping was performed only in the 60 extended FCHL families using at least 50,000 simulations. The segregating haplotype was 1-1 (1 indicates the common allele) in all gamete competition analyses above:

	FCHL all	TG all	FCHL men	TG men
Multi-HHRR	ns (ns)	0.05 (ns)	0.009 (ns)	0.00003 (0.003)
Gamete competition	0.00002 (0.005)	0.00006 (0.008)	0.0004 (0.04)	0.0000009 (0.004)
asymptotic value	p-			
Gamete competition	0.00004	0.00006	0.0004	0.00001
(Gene dropping)				
empirical p-value				

5 SUPPLEMENTARY TABLE 2. ASSOCIATION AND LINKAGE ANALYSES OF *TXNIP* WITH FCHL.

10 LOD indicates the maximum lod score of the parametric two-point or multipoint linkage analysis using the MLINK program and a dominant mode of inheritance (recombination fraction is given in parentheses); ASP indicates the lod score obtained in the affected sib-pair analysis; GAMETE indicates the p-values obtained in the Gamete competition analysis; HHRR and multi-HHRR the p-values obtained in the haplotype-based haplotype relative risk analysis; and HBAT the p-value for the test between the *TXNIP* haplotypes and the FCHL trait. Ns indicates non-significant. For the TG trait, the corresponding p-values for all association analyses remained non-significant, and both two- and multipoint lod scores were < 1.5. The numbering of the new SNP2 is based on the genomic sequence of the *TXNIP* region at the UCSC Genome Browser, July 2003. All of these SNPs were genotyped in the extended sample of 721 family members from 60 FCHL families.

Method	Analysis of single SNPs				Analysis of combined SNPs
	SNP1	SNP2	SNP3	SNP4	SNP1-2-3-4
	rs223656 7	-1273 bp C- >T	rs9245	rs7211	
Linkage					
LOD	0.4 (0.14)	0.3 (0.12)	0.3 (0.20)	0.6 (0.10)	1.9 (0.11)
ASP	0.3	0.3	0.6	0.2	
Family-based					
Association					
GAMETE	ns	ns	ns	ns	ns

HHRR	ns	ns	ns	ns	ns
HBAT					ns
Heterozygosity	0.11	0.10	0.11	0.12	

5

The *usf1s1* and *usf1s2* provided evidence for linkage in the 42 FCHL families with maximum lod scores of 3.5 and 2.0 for FCHL, and 3.7 and 2.0 for TGs. Combined analysis of these SNPs also provided some evidence for association with the gamete competition test for both FCHL ($p=0.005$) and TGs ($p=0.008$) (Table 1), although the results of individual SNPs were non-significant. We also observed a difference in the allele frequencies between unaffected and affected men, especially with the TG trait. The frequency of minor allele of *usf1s1* was 22.0% in TG-affected males and 40% in the unaffected male family members. Since these affected and unaffected family members represent non-independent groups of males, we tested *usf1s1* and *usf1s2* in TG-affected men using the family-based association method, HHRR, and the gamete competition test: p -values of 0.01 and 0.02 were obtained in the HHRR analysis and 0.008 and 0.02 in the gamete competition test of the 42 nuclear FCHL families (Table 2). The combined analysis of these SNPs yielded a p -value of 0.003 in the HHRR test and 0.004 in the gamete competition test for TGs in men (Table 1).

TABLE 2. ASSOCIATION ANALYSES OF INDIVIDUAL SNPs FOR THE *JAM1-USF1* REGION FOR TGs AND FCHL IN MEN.

All results represent p -values, ns indicates non-significant, HHRR haplotype-based haplotype relative risk test, and Gamete gamete competition test. LD cluster number in the last column indicates the clusters of SNPs showing strong intermarker LD ($p \leq 0.00002$) in the male probands with high TGs ($>90^{\text{th}}$ age-sex percentile), i.e. the SNPs carrying the same cluster number are in strong pairwise LD. SNPs indicated in bold were genotyped in the 60 extended FCHL families, and the values in parentheses

5 were obtained for these SNPs in the 42 nuclear FCHL families. All other results were obtained in the 42 nuclear FCHL families.

SNP	rs number	Distance (in bp)	Heterozygosity/Rare allele frequency in all family members	TGs HHRR	TGs Gamete	FCHL HHRR	FCHL Gamete	LD cluster (I-V)
jam1s1	rs836	1361	0.41/0.28	0.03	0.009	ns	0.03	I
jam1s2	rs790056	1561	0.36/0.24	ns	0.03	ns	ns	II
jam1s3	rs790055	25608	0.35/0.23	ns	ns	ns	ns	II
jam1s4	new	10572	0.38/0.26	0.06	0.04	ns	ns	I
jam1s5	rs4339888	1246	0.43/0.31	0.02	0.003	ns	0.09	I
jam1s6	rs3766383	951	0.25/0.15	ns	ns	ns	ns	III
usf1s1	rs3737787	1239	0.45/0.34	0.0009 (0.01)	0.00001 (0.008)	0.04 (ns)	0.05 (ns)	I
usf1s2	rs2073658	12	0.44/0.33	0.002 (0.02)	0.00006 (0.02)	0.04 (ns)	ns (ns)	I
usf1s3	rs2516841	17	0.40/0.28	ns	ns	ns	ns	II
usf1s4	rs2073657	526	0.48/0.41	ns	ns	ns	ns	IV
usf1s5	rs2516840	1443	0.41/0.29	ns	ns	ns	ns	II
usf1s6	rs2073653	361	0.25/0.14	ns	0.08	ns	ns	III
USF1 S7	rs2516839	1249	0.47/0.39	ns (ns)	0.04 (ns)	ns (ns)	ns (ns)	IV
usf1s8	rs2516838	279	0.40/0.28	0.01 (0.05)	0.05 (0.03)	ns (ns)	ns (ns)	V
usf1s9	rs1556259		0.23/0.13	ns	ns	ns	ns	III

5 SUPPLEMENTARY TABLE 3. VARIANTS IDENTIFIED BY SEQUENCING THE *USF1* GENE IN THE 31 FCHL PROBANDS OF THE ORIGINAL LINKAGE STUDY³.

Location	rs number	Rare frequencies (in 31 samples)	allele	Information on LD (in 31 samples)	Specifics
-2167	New	0.02			T/C
-2022	New	0.05			A/C
-802	New	0.03			C/G
Exon 1	rs2516837	0.44		In full LD with rs2516839 rs2774273	Not translated region
INTRON 1 = usf1s9	rs1556259	0.19			
INTRON 1 = usf1s8	rs2516838	0.29			
Intron 1	rs1556260	0.16		In full LD with SNPs in 1125 bp and 1416 bp; 30/31 samples in LD with rs1556259	
Intron 1	rs2774273	0.44		In full LD with rs2516839 and rs2516837	
Intron 1 / 1125 bp	New	0.16		In full LD with SNP 1416 bp; 30/31 samples in LD with rs1556259	C/T
Intron 1 / 1416 bp	New	0.16		In full LD with the SNP in 1125 bp; 30/31 samples in LD with rs1556259	A/G
EXON 2 = usf1s7	rs2516839	0.44			Not translated region
INTRON 2 = usf1s6	rs2073653	0.11			
Intron 3	rs2073655	0.23		In full LD with rs2073658	
Intron 5	rs2774276	0.27		29/31 in LD with rs2516840	
Intron 6	rs2073656	0.23		In full LD with	

				rs2073658
INTRON 6	rs2516840	0.32		
= usf1s5				
Intron 6 / 3411 bp	New	0.05		C/T
Intron 6 / 3519 bp	New	0.05		C/T
INTRON 7	rs2073657	0.47		In AluSx
= usf1s4				
INTRON 7	rs2516841	0.31		In AluSx
= usf1s3				
INTRON 7	rs2073658	0.23		
= usf1s2				
Intron 9 / 4445 bp	New	0.03		A/G
EXON 11	rs3737787	0.24		Not translated region
= usf1s1				

5 Underlined variants were genotyped in the FCHL families. For these SNPs, the numbers usf1s1-s9, used in the text and Tables 1-3, are also shown; New indicates that the SNP was not found in the SNP databases. The numbering of the new SNPs is based on the genomic sequence of *USF1* at the UCSC Genome Browser, July 2003 (refGene_NM_007122).

10 Next, we genotyped these two associated SNPs, usf1s1 and usf1s2, in the larger study sample of 60 extended FCHL families. Furthermore, 12 additional SNPs were genotyped for the *USF1* region (Table 2, Fig. 1). Of the 23 SNPs identified by sequencing, we genotyped all the SNPs that were not in strong LD in 31 probands, excluding six rare SNPs present in three or fewer individuals (Supplementary Table

15 2). A total of four *USF1* SNPs were genotyped in the 60 extended families due to their promising results in the nuclear study sample and/or LD pattern (Table 2). When genotyped in the 60 extended FCHL families, the two individual SNPs, usf1s1 and usf1s2, yielded p-values of 0.0009 and 0.002 in the HHRR test as well as 0.00001 and 0.0006 in the gamete competition test for TGs in men (Table 2). The

20 common allele of both SNPs was more frequently transmitted to the affected individuals in both tests and with both the FCHL and TG traits. The asymptotic p-values of the combined analyses of these two SNPs were 0.00003 in the HHRR and

- 5 0.0000009 in the combined gamete competition test for TGs in men (Table 1). The segregating haplotype was 1-1 (1 indicating the common allele). For all TG-affected family members, the combined analysis also produced evidence of association with p-values of 0.05 in the HHRR analysis and 0.00006 in the gamete competition test, again with the segregating haplotype of 1-1 (Table 1).
- 10 To confirm that the gamete competition results are indeed significant and not biased by such contributors as sparse data, we calculated empirical p-values for all gamete compete analyses involving multiple SNPs (Table 1) using gene dropping with at least 50,000 simulations (see Methods). The obtained empirical p-values were in very good agreement with the asymptotic p-values of the gamete competition
- 15 analyses (Table 1), indicating that the observed results do not represent artifacts of asymptotic approximations with sparse data.

After genotyping a total of 15 SNPs in the *USF1* region, we identified a pattern of association and LD reaching at least 46 kb in men with high TGs and extending from the centromeric junctional adhesion molecule 1 (*JAM1*) gene to the *USF1* gene

20 (Fig. 1 and Table 2): in addition to *usf1s1* and *usf1s2*, three other SNPs, *jam1s1*, *jam1s4*, and *jam1s5*, also showed evidence for association in the 42 nuclear FCHL families for high TGs in men (Table 2). These three SNPs were in strong LD with the *usf1s1* and *usf1s2* ($p < 0.00002$). The LD pattern, tested by the Genepop program, for SNPs in the *JAM1-USF1* region is shown in Table 2. In addition to these five

25 SNPs, one SNP (*usf1s8*) in intron 1 of *USF1*, showed some evidence for association as well (Table 2). This SNP was not in LD with any of the 14 other SNPs (Table 2).

In all affected family members, using both FCHL and TG traits, the evidence for association was restricted to the *usf1s1* and *usf1s2* (Table 1) within the *USF1* gene.

30 The rest of the 13 SNPs genotyped for the *JAM1-USF1* region did not provide significant evidence for association. However, we observed that two additional *USF1* SNPs among those 23 SNPs identified by sequencing, rs2073655 in intron 3 and rs2073656 in intron 6, were also in full LD with the associated *usf1s2* in 31 FCHL probands and are likely to extend the FCHL-associated region to intron 3 of

35 *USF1*. No association was obtained with SNPs residing outside the *JAM1-USF1* region (Supplementary Table 1). In conclusion, evidence for association and LD was restricted to a 1239 bp region within the *USF1* gene in all affected individuals of

- 5 FCHL families but extended at least 46 kb within the *JAM1-USF1* region in men with high TGs (Tables 2-3, Fig. 1).

The combination of the *usf1s1-usf1s2* SNPs, resulting in the significant haplotypes for FCHL and TGs, was also tested with three additional qualitative lipid traits: high apolipoprotein B (apoB), high TC and small low-density lipoprotein (LDL) peak
 10 particle size. For apoB, p-values of 0.00003 and 0.0007 were obtained for all affected individuals and for affected men for the susceptibility haplotype 1 -1 in the gamete competition analysis. For TC, the p-values were 0.0001 and 0.007; and for LDL peak particle size, 0.002 and 0.01, respectively. These results together with the results obtained for FCHL suggest that the underlying gene is not affecting TGs
 15 alone but also the complex FCHL phenotype.

EXAMPLE 3: HAPLOTYPE ANALYSES OF THE *JAM1-USF1* GENE REGION

Using the HBAT program we obtained evidence for shared haplotypes in the region of *usf1s1* and *usf1s2* (Table 3). This observation was supported by multipoint HHRR
 20 analyses (Table 3). For the haplotype 1-1 (1 indicating the common allele) a p-value of 0.0007 was obtained using the -o option.

TABLE 3. HAPLOTYPE ANALYSES IN TG-AFFECTED MEN USING THE HBAT PROGRAM (THE MULTILOCUS GENO-PDT AND MULTI-HHRR RESULTS ARE GIVEN BELOW FOR COMPARISON).

25 The inter-SNP distances and corresponding rs numbers for the SNPs *jam1s4-s6* and *usf1s1-s5* are shown in Table 2; 1 indicates the common allele; and ns non-significant. The p-value of the HBAT program indicates the probability that the particular haplotype is transmitted to the affected
 30 individuals using the option -o (optimize offset) or option -e (empirical test). Multilocus geno-PDT indicates a genotype-based association test for general pedigrees. The multi-HHRR analysis is testing the hypothesis of homogeneity of marker allele distributions between transmitted and non-transmitted alleles of the SNPs.

Test	Haplotype of SNPs: Jam1s4-6 - usf1s1-2	Haplotype of SNPs: usf1s1-2	Haplotype of SNPs: usf1s1-5
HBAT -o	P = 0.03 (haplotype 1-1-1-1-1)	P = 0.0007 (haplotype 1-1)	P = ns (0.07) (haplotype 1-1-1-1-1)
		P = 0.004 for the protective haplotype 2-2, significantly less transmitted to the affected subjects	
HBAT -e	P = 0.009 (haplotype 1-1-1-1-1)	P = 0.02 (haplotype 1-1)	P = ns (0.2) (haplotype 1-1-1-1-1)
Multi-locus geno- PDT	P = 0.02	P = 0.002	P = ns (0.7)
Multi- HHRR	P = 0.0002	P = 0.00003	P = 0.04

5

This option measures not only preferential transmission of the susceptibility haplotype to affecteds but also less preferential transmissions to unaffecteds, making it useful here since in these extended families the unaffecteds also contain important information. The results of the HBAT -e option, a test of association given linkage, are also shown in Table 3. Since this test statistics implicitly conditions on linkage information, it is less powerful and leads to reduced p-values. However, this test together with the results of the HHRR analyses allow us to conclude that the 1-1 haplotype is associated with the phenotype (Table 3). Furthermore, haplotype 2-2 was significantly less transmitted to the affected subjects ($p=0.004$), suggesting a protective role for this allele. These results were further supported by a genotype-based association test for general pedigrees, the genotype-PDT, which provided evidence for association (Table 3), as well as by the gamete competition analyses (Table 1), where the same haplotype 1-1 was segregating to the affected individuals with both FCHL and TG traits.

5 EXAMPLE 4: EXPRESSION PROFILES OF FAT BIOPSIES AND INITIAL FUNCTIONAL ANALYSIS

We investigated whether the gene expression profiles of fat biopsies from six affected FCHL family members carrying the susceptibility haplotype 1-1, constructed by the SNPs *usf1s1* and *usf1s2*, revealed differences when compared to four
10 affected FCHL family members homozygous for the putative protective haplotype, 2-2 (see above), using the Affymetrix, HGU133A probe array. We also specifically investigated whether *USF1* is expressed in fat tissue because it is not sufficiently represented on the Affymetrix HGU133A chip. Using RT-PCR the *USF1* was found to be expressed in the fat biopsy samples (data not shown). Quantitative real-time
15 PCR was also performed to determine the relative expression levels of *USF1* in adipose tissue in the affected FCHL family members carrying the risk haplotype and affected members not carrying the risk haplotype. No detectable differences in *USF1* expression levels could be observed, suggesting that the potential functional significance of the FCHL associated allele of the *USF1* is not delivered via a direct
20 effect on the steady state transcript level in adipose tissue.

Due to the limited number of samples available, statistical power to detect differences in gene expression between the haplotype groups was not considered sufficient. As an alternative, we therefore defined cut-off thresholds (see Methods) to discriminate between significant differences and differences attributable to
25 technical or biological noise in the experimental procedures. Using these criteria, we identified 25 genes that appeared up-regulated and 73 genes down-regulated in the susceptibility haplotype carriers (the complete lists will be available at our website, while the raw data can be accessed through the Gene Expression Omnibus at NCBI using the GEO accession GSE590). To lend biological relevance to these findings,
30 lists of differentially expressed genes were examined for over-representation of functional classes, as defined by the gene ontology (GO) consortium, using the Expression Analysis Systematic Explorer (EASE) tool. Only three classes were found to be statistically significantly over-represented among the up-regulated genes (Fig. 2), primarily implicating genes involved in fat metabolism. Among the
35 down-regulated genes, a prominent down-regulation of immune-response genes was observed (Fig. 2). The complete results from the EASE analysis, including the corresponding EASE scores (p-values) and lists of genes in the significant (=p-value<0.05) functional categories, are given in the Supplementary Table 3a-b.

5 Next we investigated the genomic sequence flanking the haplotype 1-1, and identified a 60-bp sequence element found in 91 human genes as follows: The SNP usf1s2, forming part of the haplotype 1-1, resides adjacent (8 bp) to a 306-bp AluSx repeat. Two parts (2-61 bp and 137-196 bp) of this AluSx repeat show sequence similarity with the mouse B1 repeat (Fig. 3a). When blasted against the mouse
 10 sequence databases, these two parts of the AluSx sequence identify numerous mouse ESTs, due to the B1 element located in the untranslated region of the mouse mRNA. When blasted against human sequence databases, 91 human genes, including *USF1*, have this 60-bp part of AluSx either on the coding strand (43 genes) or on the opposite strand (48 genes). The 60-bp part is highly conserved
 15 from human to worm since it was found in pufferfish and *Caenorhabditis elegans* but not in *Drosophila melanogaster* or in *Saccharomyces cerevisiae*. A complete list of the 91 human genes as well as their individual p-values and identity percentages (between 83-98%) are given in Supplementary Table 4. Analysis of domain annotation of the 91 genes indicates enrichment of domains involved in protein
 20 modification (n=16) and domains related to nucleic acids (n=35). This observation was also supported by the available annotations about biological process, where majority of the genes were involved in nucleic acid metabolism (n=18), as well as in transcription and signal transduction (n=33).

To obtain some evidence for the functional significance of this conserved 60-bp
 25 DNA element, we produced a 268-bp long construct containing the critical 60-bp sequence as well as the usf1s2 SNP region and tested its regulatory function in vitro using the SEAP reporter system (Fig. 3b). The genomic DNAs from one homozygous susceptibility carrier (haplotype 1-1) and one homozygous non-carrier (2-2) were cloned in front of the SEAP reporter gene in two orientations. The effect
 30 on the transcription of the reporter gene was implicated in the forward orientation in both constructs, whereas the reverse orientation resulted in the transcription efficiency comparable to the negative control (Fig. 3b).

5 SUPPLEMENTARY TABLE 4A. RESULTS FROM ANALYSIS OF LISTS WITH DIFFERENTIALLY EXPRESSED GENES BETWEEN THE HAPLOTYPE CARRIERS AND NON-CARRIERS FOR OVER-REPRESENTATION OF FUNCTIONAL CATEGORIES USING THE EASE TOOL²⁷. THIS SUPPLEMENTARY TABLE 4A-B WILL BE SHOWN AT OUR WEB SITE. PLEASE SEE FIGURE 2 FOR THE GRAPHICAL DISTRIBUTION OF THESE GENES ACCORDING TO THE FUNCTIONAL CATEGORY.

Functional category ¹	LH	LT	PH	PT	EASE score (p-value)
UP-REGULATED GENES					
fatty acid metabolism	3	16	90	7689	0.0129
lipid metabolism	4	16	359	7689	0.0302
macromolecule catabolism	4	16	395	7689	0.0386
carboxylic acid metabolism	3	16	230	7689	0.0724
organic acid metabolism	3	16	232	7689	0.0735
cell motility	3	16	253	7689	0.0855
Catabolism	4	16	554	7689	0.0885
proteolysis and peptidolysis	3	16	368	7689	0.159
protein catabolism	3	16	374	7689	0.164
Metabolism	11	16	4163	7689	0.239
cell proliferation	3	16	782	7689	0.46

Physiological processes	14	16	6379	7689	0.516
cell growth and/or maintenance	6	16	2389	7689	0.521
protein metabolism	4	16	1512	7689	0.589
cellular process	9	16	4297	7689	0.679
cell communication	4	16	2238	7689	0.858
nucleobase, nucleoside, nucleotide and nucleic acid metabolism	3	16	1716	7689	0.88
Functional category ¹					
	LH	LT	PH	PT	EASE score (p-value)
Down-regulated genes					
immune response	16	60	560	7689	0.0000141
response to pest/pathogen/parasite	13	60	379	7689	0.0000236
response to biotic stimulus	17	60	674	7689	0.0000309
defense response	16	60	616	7689	0.0000435
response to wounding	9	60	222	7689	0.000265
response to stress	14	60	632	7689	0.000811

inflammatory response	7	60	149	7689	0.000926
innate immune response	7	60	151	7689	0.000992
response to external stimulus	17	60	992	7689	0.00256
Catabolism	12	60	554	7689	0.00288
colony morphology	3	60	26	7689	0.0167
invasive growth	3	60	26	7689	0.0167
cytosolic calcium ion concentration elevation	3	60	32	7689	0.0248
cellular morphogenesis	3	60	34	7689	0.0277
cell adhesion	8	60	390	7689	0.0287
macromolecule catabolism	8	60	395	7689	0.0305
lipid catabolism	3	60	50	7689	0.0561
proteolysis and peptidolysis	7	60	368	7689	0.0617
protein catabolism	7	60	374	7689	0.0657
G-protein signaling, coupled to IP3 second messenger (phospholipase C activating)	3	60	66	7689	0.0909
Endocytosis	3	60	72	7689	0.105

cellular defense response	3	60	77	7689	0.118
lipid metabolism	6	60	359	7689	0.14
Chemotaxis	3	60	90	7689	0.151
Taxis	3	60	90	7689	0.151
Antimicrobial humoral response	3	60	92	7689	0.157
humoral defense mechanism (sensu Invertebrata)	3	60	92	7689	0.157
antimicrobial humoral response (sensu Invertebrata)	3	60	92	7689	0.157
vesicle-mediated transport	4	60	214	7689	0.226
cell-cell adhesion	3	60	136	7689	0.28
response to chemical substance	3	60	141	7689	0.295
alcohol metabolism	3	60	149	7689	0.317
humoral immune response	3	60	152	7689	0.326
cell surface receptor linked signal transduction	8	60	739	7689	0.338
cell communication	20	60	2238	7689	0.345
signal transduction	16	60	1785	7689	0.392

cell death	4	60	313	7689	0.433
Death	4	60	316	7689	0.439
Physiological processes	51	60	6379	7689	0.439
G-protein coupled receptor protein signaling pathway	5	60	457	7689	0.469
metal ion transport	3	60	216	7689	0.497
protein metabolism	13	60	1512	7689	0.5
phosphate metabolism	5	60	487	7689	0.519
phosphorus metabolism	5	60	487	7689	0.519
Transport	10	60	1144	7689	0.524
Development	10	60	1165	7689	0.547
cellular process	34	60	4297	7689	0.551
morphogenesis	6	60	669	7689	0.592
Carbohydrate metabolism	3	60	261	7689	0.6
ion transport	4	60	410	7689	0.616
cation transport	3	60	288	7689	0.655

Apoptosis	3	60	289	7689	0.656
Programmed cell death	3	60	290	7689	0.658
cell organization and biogenesis	4	60	437	7689	0.66
intracellular signaling cascade	5	60	596	7689	0.681
cell growth and/or maintenance	18	60	2389	7689	0.693
Metabolism	31	60	4163	7689	0.74
protein amino acid phosphorylation	3	60	365	7689	0.778
response to abiotic stimulus	3	60	389	7689	0.807
phosphorylation	3	60	393	7689	0.812
organogenesis	4	60	637	7689	0.878
protein modification	4	60	682	7689	0.905
cell proliferation	3	60	782	7689	0.987
regulation of transcription, DNA-dependent	3	60	974	7689	0.997
regulation of transcription	3	60	979	7689	0.997
transcription, DNA-dependent	3	60	1085	7689	0.999
Transcription	3	60	1112	7689	0.999

nucleobase, nucleoside, nucleotide and nucleic acid metabolism	5	60	1716	7689	1
--	---	----	------	------	---

5 ¹According to the gene ontology (GO) classification biological process⁴¹. Abbreviations: LH - list hits, LT - list total, PH - population hits, PT - population total, and EASE - Expression Analysis Systematic Explorer⁴². The complete lists of genes in each functional category will be presented at our web site.

10 Supplementary Table 4b. Lists of genes in the significant (= EASE p-value <0.05) functional categories in Table 3a above. This supplementary Table 4a-b will be shown at our web site.

Up-regulated genes

Fatty acid metabolism

15

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
200832_s_at	6319	stearoyl-CoA desaturase (delta-9-desaturase)	biological_process	endoplasmic reticulum; fatty acid biosynthesis; integral to membrane; iron ion binding; oxidoreductase activity; stearoyl-CoA desaturase activity

206930_at	10249	glycine-N-acyltransferase	biological_process	acyl-CoA metabolism; acyltransferase activity; mitochondrion; response to toxin
209600_s_at	51	acyl-Coenzyme A oxidase 1, palmitoyl	biological_process	acyl-CoA oxidase activity; electron donor activity; electron transport; energy pathways; fatty acid beta-oxidation; oxidoreductase activity; peroxisome; prostaglandin metabolism

Lipid metabolism

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
200832_s_at	6319	stearoyl-CoA desaturase (delta-9-desaturase)	biological_process	endoplasmic reticulum; fatty acid biosynthesis; integral to membrane; iron ion binding; oxidoreductase activity; stearoyl-CoA desaturase activity
202118_s_at	8895	copine III	biological_process	calcium-dependent phospholipid binding; cell adhesion molecule activity; lipid metabolism; transporter activity; vesicle-mediated transport
206930_at	10249	glycine-N-	biological_process	acyl-CoA metabolism; acyltransferase activity; mitochondrion; response to

acyltransferase		toxin	
209600_s_51 at	acyl-Coenzyme A oxidase 1, palmitoyl	biological_process	acyl-CoA oxidase activity; electron donor activity; electron transport; energy pathways; fatty acid beta-oxidation; oxidoreductase activity; peroxisome; prostaglandin metabolism

5

Macromolecule catabolism

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
202581_at	3304	heat shock 70kDa protein 1B	biological_process	ATP binding; cytoplasm; heat shock protein activity; mRNA catabolism; nucleus
204844_at	2028	glutamyl aminopeptidase (aminopeptidase A)	biological_process	cell proliferation; cell-cell signaling; glutamyl aminopeptidase activity; hydrolase activity; integral to plasma membrane; membrane alanyl aminopeptidase activity; metallopeptidase activity; proteolysis and peptidolysis; zinc ion binding
209788_s_51752 at		type 1 tumor necrosis factor receptor shedding	biological_process	aminopeptidase activity; membrane alanyl aminopeptidase activity; metallopeptidase activity; proteolysis and peptidolysis; zinc ion binding

aminopeptidase regulator			
215271_at	63923	tenascin N	biological_process carboxypeptidase A activity; cell growth; cell migration; cellular_component unknown; molecular_function unknown; proteolysis and peptidolysis

Down-regulated genes

Immune response

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201422_at	10437	interferon, inducible protein 30	gamma- biological_process	extracellular; immune response; lysosome; oxidoreductase activity
201952_at	214	activated leukocyte adhesion molecule	cell biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_ at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway

(mac-1) beta subunit)				
202901_x_ at	1520	cathepsin S	biological_process	cathepsin S activity; hydrolase activity; immune response; lysosome; proteolysis and peptidolysis
203104_at	1436	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	biological_process	ATP binding; antimicrobial humoral response (sensu Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203382_s_ at	348	apolipoprotein E	biological_process	cholesterol metabolism; circulation; development; heparin binding; immune response; lipid binding; lipid metabolism; lipid transport; lipid transporter activity; receptor binding
203650_at	10544	protein C receptor, endothelial (EPCR)	biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204122_at	7305	TYRO protein tyrosine kinase binding protein	biological_process	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_ at	240	arachidonate lipoxygenase	5- biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell

206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding	signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
209906_at	719	complement component 3a receptor 1	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction	
211530_x_at	3135	HLA-G histocompatibility antigen, class I, G	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite	
211799_x_at	3107	major histocompatibility complex, class I, C	biological_process	MHC class II receptor activity; class I major histocompatibility complex antigen; immune response; integral to membrane	
213975_s_at	4069	lysozyme (renal amyloidosis)	biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity	
217028_at	7852	chemokine (C-X-C motif)	biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor	

receptor 4	activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity
------------	---

Response to pest/pathogen/parasite

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201850_at	822	capping protein (actin filament), gelsolin-like	biological_process	F-actin capping protein complex; actin binding; barbed-end actin capping activity; nucleus; protein complex assembly; response to pest/pathogen/parasite
201952_at	214	activated leukocyte cell adhesion molecule	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) beta subunit)	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
203104_at	1436	colony stimulating factor 1	biological_process	ATP binding; antimicrobial humoral response (sensu

		receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog		Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203650_at	10544	protein C receptor, endothelial (EPCR)	biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204122_at	7305	TYRO protein tyrosine kinase binding protein	biological_process	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_at	240	arachidonate lipooxygenase	5- biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense

211530_x_at	3135	HLA-G antigen, class I, G	histocompatibility	biological_process	response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
213975_s_at	4069	lysozyme amyloidosis)	(renal	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite
217028_at	7852	chemokine receptor 4	(C-X-C motif	biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
					C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Response to biotic stimulus

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201422_at	10437	interferon, gamma-inducible	biological_process	extracellular; immune response; lysosome; oxidoreductase activity

protein 30	s		
201850_at	822	capping protein (actin filament), gelsolin-like	biological_processes F-actin capping protein complex; actin binding; barbed-end actin capping activity; nucleus; protein complex assembly; response to pest/pathogen/parasite
201952_at	214	activated leukocyte adhesion molecule	biological_processes antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) (beta subunit)	biological_processes antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
202901_x_at	1520	cathepsin S	biological_processes cathepsin S activity; hydrolase activity; immune response; lysosome; proteolysis and peptidolysis
203104_at	1436	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	biological_processes ATP binding; antimicrobial humoral response (sensu Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203382_s_at	348	apolipoprotein E	biological_processes cholesterol metabolism; circulation; development; heparin binding; immune response; lipid binding; lipid metabolism; lipid transport; lipid transporter activity; receptor binding

203650_at	10544	protein C receptor, endothelial (EPCR)	biological_processes	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204122_at	7305	TYRO protein tyrosine kinase binding protein	biological_processes	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
04446_s_at	240	arachidonate lipoxigenase	5-biological_processes	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_processes	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	biological_processes	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1	biological_processes	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like

211530_x_at	3135	HLA-G antigen, class I, G	histocompatibility	biological_processes	receptor activity; smooth muscle contraction
					MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite
211799_x_at	3107	major histocompatibility complex, class I, C	histocompatibility	biological_processes	MHC class II receptor activity; class I major histocompatibility complex antigen; immune response; integral to membrane
213975_s_at	4069	lysozyme amyloidosis)	(renal	biological_processes	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
217028_at	7852	chemokine (C-X-C motif) receptor 4	(C-X-C motif)	biological_processes	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Defense response

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
--------	-----------	----------	----------------	---------------------------

201422_at	10437	interferon, inducible protein 30	gamma-cell	biological_process	extracellular; immune response; lysosome; oxidoreductase activity
201952_at	214	activated leukocyte adhesion molecule	cell	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) (beta subunit)		biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
202901_x_at	1520	cathepsin S		biological_process	cathepsin S activity; hydrolase activity; immune response; lysosome; proteolysis and peptidolysis
203104_at	1436	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog		biological_process	ATP binding; antimicrobial humoral response (sensu Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203382_s_at	348	apolipoprotein E		biological_process	cholesterol metabolism; circulation; development; heparin binding; immune response; lipid binding; lipid metabolism; lipid transport; lipid transporter activity; receptor binding
203650_at	10544	protein C receptor		biological_process	blood coagulation; inflammatory response; integral to plasma

204122_at	7305	TYRO	protein tyrosine kinase binding protein	biological_process	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_at	240	arachidonate lipoxigenase	5-	biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine receptor 1	(C-C motif)	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)		biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1		biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
211530_x_at	3135	HLA-G antigen, class I, G	histocompatibility antigen, class I, G	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC

211799_x_ at	3107	major complex, class I, C	histocompatibility biological_process	class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite
213975_s_ at	4069	lysozyme amyloidosis)	(renal biological_process	MHC class II receptor activity; class I major histocompatibility complex antigen; immune response; integral to membrane
217028_at	7852	chemokine receptor 4	(C-X-C motif) biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
				C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Response to wounding

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
203650_at	10544	protein C endothelial (EPCR)	receptor, biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity

204122_at	7305	TYRO protein tyrosine kinase binding protein	biological_process	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_at	240	arachidonate lipoxigenase	5- biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating acetylhydrolase, plasma)	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
211530_x_at	3135	HLA-G histocompatibility antigen, class I, G	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite

213975_s_at	4069	lysozyme amyloidosis)	(renal biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
217028_at	7852	chemokine motif) receptor 4	(C-X-C biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

5

Response to stress

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201739_at	6446	serum/glucocorticoid regulated kinase	biological_process	ATP binding; apoptosis; cAMP-dependent protein kinase activity; protein amino acid phosphorylation; protein kinase CK2 activity; protein serine/threonine kinase activity; response to stress; sodium ion transport; transferase activity
201850_at	822	capping protein filament), gelsolin-like	(actin biological_process	F-actin capping protein complex; actin binding; barbed-end actin capping activity; nucleus; protein complex assembly; response to pest/pathogen/parasite

201952_at	214	activated leukocyte adhesion molecule	cell	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) beta subunit)		biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
203104_at	1436	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog		biological_process	ATP binding; antimicrobial humoral response (sensu Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203650_at	10544	protein C receptor, endothelial (EPCR)		biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204122_at	7305	TYRO protein tyrosine kinase binding protein		biological_process	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_at	240	arachidonate 5-lipoxygenase		biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1		biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration

206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	biological_process	elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
209906_at	719	complement component 3a receptor 1	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
211530_x_at	3135	HLA-G histocompatibility antigen, class I, G	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
213975_s_at	4069	lysozyme (renal amyloidosis)	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite
217028_at	7852	chemokine (C-X-C motif) receptor 4	biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
				C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Response to stress

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201739_at	6446	serum/glucocorticoid regulated kinase	biological_process	ATP binding; apoptosis; cAMP-dependent protein kinase activity; protein amino acid phosphorylation; protein kinase CK2 activity; protein serine/threonine kinase activity; response to stress; sodium ion transport; transferase activity
201850_at	822	capping protein (actin filament), gelsolin-like	biological_process	F-actin capping protein complex; actin binding; barbed-end actin capping activity; nucleus; protein complex assembly; response to pest/pathogen/parasite
201952_at	214	activated leukocyte adhesion molecule	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) beta subunit)	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
203104_at	1436	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma	biological_process	ATP binding; antimicrobial humoral response (sensu Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor

		viral (v-fms) oncogene homolog	receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203650_at	10544	protein C receptor, endothelial (EPCR)	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204122_at	7305	TYRO protein tyrosine kinase binding protein	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_at	240	arachidonate 5-lipoxygenase	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to

211530_x_at	3135	HLA-G antigen, class I, G	histocompatibility	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite	plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
213975_s_at	4069	lysozyme (renal amyloidosis)		biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity	
217028_at	7852	chemokine (C-X-C motif) receptor 4		biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity	

Inflammatory response

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
203650_at	10544	protein C receptor, endothelial (EPCR)	biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity

204446_s_at	240	arachidonate lipooxygenase	5- biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase group VII (platelet-activating acetylhydrolase, plasma)	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
213975_s_at	4069	lysozyme (renal amyloidosis)	biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
217028_at	7852	chemokine (C-X-C motif) receptor 4	biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of

MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Innate immune response

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
203650_at	10544	protein C receptor, endothelial (EPCR)	biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204446_s_at	240	arachidonate lipooxygenase	5- biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase,	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding

plasma)			
209906_at	719	complement component receptor 1	biological_process 3a C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
213975_s_at	4069	lysozyme amyloidosis)	(renal biological_process carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
217028_at	7852	chemokine (C-X-C motif) receptor 4	biological_process C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Response to external stimulus

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201422_at	10437	interferon,	gamma- biological_process	extracellular; immune response; lysosome; oxidoreductase

inducible protein 30		activity	
201850_at	822	capping protein (actin filament), gelsolin-like	biological_process F-actin capping protein complex; actin binding; barbed-end actin capping activity; nucleus; protein complex assembly; response to pest/pathogen/parasite
201952_at	214	activated leukocyte adhesion molecule	biological_process antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) (beta subunit)	biological_process antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
202901_x_at	1520	cathepsin S	biological_process cathepsin S activity; hydrolase activity; immune response; lysosome; proteolysis and peptidolysis
203104_at	1436	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	biological_process ATP binding; antimicrobial humoral response (sensu Invertebrata); cell proliferation; development; integral to plasma membrane; macrophage colony stimulating factor receptor activity; protein amino acid phosphorylation; receptor activity; signal transduction; transferase activity; transmembrane receptor protein tyrosine kinase signaling pathway
203382_s_at	348	apolipoprotein E	biological_process cholesterol metabolism; circulation; development; heparin binding; immune response; lipid binding; lipid metabolism; lipid transport; lipid transporter activity; receptor binding

203650_at	10544	protein C receptor, endothelial (EPCR)	biological_process	blood coagulation; inflammatory response; integral to plasma membrane; receptor activity
204122_at	7305	TYRO protein tyrosine kinase binding protein	biological_process	cellular defense response; integral to plasma membrane; intracellular signaling cascade; receptor signaling protein activity
204446_s_at	240	arachidonate lipoxigenase	5- biological_process	arachidonate 5-lipoxygenase activity; electron transport; inflammatory response; iron ion binding; leukotriene biosynthesis; lipoxygenase activity; oxidoreductase activity
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
209906_at	719	complement component 3a receptor 1	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like

211530_x_at	3135	HLA-G antigen, class I, G	histocompatibility	biological_process	receptor activity; smooth muscle contraction
211799_x_at	3107	major complex, class I, C	histocompatibility	biological_process	MHC class I receptor activity; antigen presentation, endogenous antigen; antigen processing, endogenous antigen via MHC class I; cellular defense response; integral to membrane; perception of pest/pathogen/parasite
213975_s_at	4069	lysozyme amyloidosis)	(renal	biological_process	MHC class II receptor activity; class I major histocompatibility complex antigen; immune response; integral to membrane
217028_at	7852	chemokine (C-X-C motif) receptor 4		biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
					C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Catabolism

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
--------	-----------	----------	----------------	---------------------------

202295_s_ at	1512	cathepsin H	biological_process	cathepsin H activity; hydrolase activity; lysosome; proteolysis and peptidolysis
202901_x_ at	1520	cathepsin S	biological_process	cathepsin S activity; hydrolase activity; immune response; lysosome; proteolysis and peptidolysis
203649_s_ at	5320	phospholipase A2, group IIA (platelets, synovial fluid)	biological_process	calcium ion binding; calcium-dependent cytosolic phospholipase A2 activity; calcium-dependent secreted phospholipase A2 activity; calcium-independent cytosolic phospholipase A2 activity; hydrolase activity; lipid catabolism; membrane
203936_s_ at	4318	matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)	biological_process	collagen catabolism; collagenase activity; extracellular matrix; extracellular space; gelatinase B activity; hydrolase activity; zinc ion binding
206214_at	7941	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)	biological_process	2-acetyl-1-alkylglycerophosphocholine esterase activity; 2-acetyl-1-alkylglycerophosphocholine esterase complex; extracellular; hydrolase activity; inflammatory response; lipid catabolism; phospholipid binding
207332_s_ at	7037	transferrin receptor (p90, CD71)	biological_process	endocytosis; endosome; extracellular; integral to plasma membrane; iron ion homeostasis; iron ion transport; peptidase activity; proteolysis and peptidolysis; receptor activity; transferrin receptor activity
213274_s_ at	1508	cathepsin B	biological_process	cathepsin B activity; hydrolase activity; intracellular; lysosome; proteolysis and peptidolysis

213510_x_at	220594	TL132 protein	biological_process	cysteine-type endopeptidase activity; ubiquitin C-terminal hydrolase activity; ubiquitin-dependent protein catabolism
213975_s_at	4069	lysozyme (renal amyloidosis)	biological_process	carbohydrate metabolism; cell wall catabolism; cytolysis; extracellular space; hydrolase activity, acting on glycosyl bonds; inflammatory response; lysin activity; lysozyme activity
214012_at	51752	type 1 tumor necrosis factor receptor shedding aminopeptidase regulator	biological_process	aminopeptidase activity; membrane alanyl aminopeptidase activity; metalloproteinase activity; proteolysis and peptidolysis; zinc ion binding
217983_s_at	8635	ribonuclease 6 precursor	biological_process	RNA catabolism; extracellular; ribonuclease activity
35820_at	2760	GM2 ganglioside activator protein	biological_process	glycolipid catabolism; glycosphingolipid metabolism; lysosome; sphingolipid activator protein activity; sphingolipid catabolism

Colony morphology

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
--------	-----------	----------	----------------	---------------------------

203186_s_6275 at	S100 calcium binding protein A4 (calcium protein, calvasculin, metastasin, murine placental homolog)	biological_process	calcium ion binding; invasive growth
205098_at	1230 chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
217028_at	7852 chemokine (C-X-C motif) receptor 4	biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Invasive growth

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
203186_s_6275 at	6275	S100 calcium binding protein A4 (calcium protein, calvasculin, metastasin, murine	biological_process	calcium ion binding; invasive growth

placental homolog)

205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
217028_at	7852	chemokine (C-X-C motif) receptor 4	biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Cystolic calcium ion concentration elevation

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
209906	719	complement	biological_process	C3a anaphylatoxin receptor activity; G-protein coupled receptor protein

at	component 3a receptor 1	signaling pathway; cell motility; cellular defense response; chemotaxis; circulation; complement component C3a receptor activity; cytosolic calcium ion concentration elevation; inflammatory response; integral to plasma membrane; phosphatidylinositol-4,5-bisphosphate hydrolysis; rhodopsin-like receptor activity; smooth muscle contraction
217028_7852 at	chemokine (C-X-C motif) receptor 4	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity

Cellular morphogenesis

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
203186_s_6275 at		S100 calcium binding protein A4 (calcium protein, calvasculin, metastasin, murine placental homolog)	biological_process	calcium ion binding; invasive growth
205098_at	1230	chemokine (C-C motif) receptor 1	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity

217028_at	7852	chemokine motif) receptor 4	(C-X-C biological_process	C-C chemokine receptor activity; C-X-C chemokine receptor activity; G-protein coupled receptor protein signaling pathway; activation of MAPK; apoptosis; chemotaxis; coreceptor activity; cytoplasm; cytosolic calcium ion concentration elevation; histogenesis and organogenesis; immune response; inflammatory response; integral to plasma membrane; invasive growth; neurogenesis; pathogenesis; response to viruses; rhodopsin-like receptor activity
-----------	------	--------------------------------	---------------------------	---

Cell adhesion

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
201952_at	214	activated leukocyte cell adhesion molecule	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion molecule activity; integral to plasma membrane; membrane fraction; receptor binding; signal transduction
202803_s_at	3689	integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) beta subunit)	biological_process	antimicrobial humoral response (sensu Invertebrata); cell adhesion; cell adhesion receptor activity; integrin complex; integrin-mediated signaling pathway
204438_at	4360	mannose receptor, C type 1	biological_process	calcium ion binding; heterophilic cell adhesion; integral to plasma membrane; mannose binding; pinocytosis; receptor activity; receptor mediated endocytosis; sugar binding

204620_s_at	1462	chondroitin proteoglycan 2 (versican)	sulfate	biological_process	calcium ion binding; cell recognition; development; extracellular matrix; heterophilic cell adhesion; hyaluronic acid binding; sugar binding
205098_at	1230	chemokine receptor 1	(C-C motif)	biological_process	C-C chemokine receptor activity; G-protein signaling, coupled to cyclic nucleotide second messenger; cell adhesion; cell-cell signaling; chemotaxis; cytosolic calcium ion concentration elevation; immune response; inflammatory response; integral to plasma membrane; invasive growth; rhodopsin-like receptor activity
205786_s_at	3684	integrin, (complement component receptor 3, alpha; also known as CD11b (p170), macrophage antigen alpha polypeptide)	alpha M	biological_process	cell adhesion; cell adhesion receptor activity; integrin complex
212014_x_at	960	CD44 antigen (homing function and Indian blood group system)		biological_process	cell adhesion receptor activity; cell-cell adhesion; cell-matrix adhesion; collagen binding; hyaluronic acid binding; integral to plasma membrane; receptor activity
216442_x_at	2335	fibronectin 1		biological_process	cell adhesion; cell adhesion molecule activity; cell motility; extracellular matrix; extracellular space; signal transduction; soluble fraction

UNIQID	LOCUSLINK	GENENAME	CLASSIFICATION	LOCUSLINK CLASSIFICATIONS
202295_s_ at	1512	cathepsin H	biological_process	cathepsin H activity; hydrolase activity; lysosome; proteolysis and peptidolysis
202901_x_ at	1520	cathepsin S	biological_process	cathepsin S activity; hydrolase activity; immune response; lysosome; proteolysis and peptidolysis
203936_s_ at	4318	matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)	biological_process	collagen catabolism; collagenase activity; extracellular matrix; extracellular space; gelatinase B activity; hydrolase activity; zinc ion binding
207332_s_ at	7037	transferrin receptor (p90, CD71)	biological_process	endocytosis; endosome; extracellular; integral to plasma membrane; iron ion homeostasis; iron ion transport; peptidase activity; proteolysis and peptidolysis; receptor activity; transferrin receptor activity
213274_s_ at	1508	cathepsin B	biological_process	cathepsin B activity; hydrolase activity; intracellular; lysosome; proteolysis and peptidolysis
213510_x_ at	220594	TL132 protein	biological_process	cysteine-type endopeptidase activity; ubiquitin C-terminal hydrolase activity; ubiquitin-dependent protein catabolism

214012_at	51752	type 1 tumor necrosis factor receptor shedding aminopeptidase regulator	biological_process	aminopeptidase activity; membrane alanyl aminopeptidase activity; metalloproteinase activity; proteolysis and peptidolysis; zinc ion binding
217983_s_ at	8635	ribonuclease 6 precursor	biological_process	RNA catabolism; extracellular; ribonuclease activity

5 The purpose of this experiment was not to solve whether the *usf1s2* SNP is directly
 causative to FCHL. More complex functional studies need to be performed before
 any conclusions of the functional significance of a single non-coding SNP can be
 drawn. However, these preliminary data combined with the across species
 conservation would imply that the DNA region flanking the susceptibility haplotype
 10 contains an element affecting transcriptional regulation. The data also suggest that
 the element is more likely to be a *Cis* acting type regulator rather than a direction-
 independent enhancer element.

EXAMPLE 5: EXPERIMENTAL SETUP - METHODS

15 The Finnish FCHL families were recruited in the Helsinki, Turku and Kuopio
 University Central Hospitals, as described earlier^{4,9}. Each subject provided a written
 informed consent prior to participating in the study. All samples were collected in
 accordance with the Helsinki declaration, and the ethics committees of the
 participating centers approved the study design. The inclusion criteria for the FCHL
 20 probands were as follows⁴: 1) serum TC and/or TGs > 90th age-sex specific Finnish
 population percentiles⁴, but if the proband had only one elevated lipid trait, a first-
 degree relative had to have the combined phenotype; 2) age > 30 years and < 55
 for males and < 65 years for females; 3) at least a 50% stenosis in one or more
 coronary arteries in coronary angiography. Exclusion criteria for the FCHL probands
 25 were type 1 DM, hepatic or renal disease, and hypothyroidism. Familial
 hypercholesterolemia was excluded from each pedigree by determining the LDL-
 receptor status of the proband by the lymphocyte culture method⁴. If the above
 mentioned criteria were fulfilled, families with at least two affected members were
 included in the study, and all the accessible family members were examined. Two
 30 traits were analyzed: FCHL and TGs. For the FCHL trait, family members were
 scored as affected according to the same diagnostic criteria as in our original
 linkage study⁴ using the Finnish age-sex specific 90th percentiles for high TC and
 high TGs, available from the web site of the National Public Health Institute, Finland.
 These ascertainment criteria are fully comparable with the original criteria¹. For
 35 analysis of TGs, family members with TG levels \geq 90th Finnish age-sex specific

5 population percentile were coded as affected. In addition to the FCHL and TG traits, the combination of the *usf1s1-usf1s2* SNPs, which resulted in the significant haplotypes for the FCHL and TG traits, was also analyzed using the apolipoprotein B (apoB), LDL peak particle size and TC traits. For apoB and TC, the 90th age-sex specific Finnish population percentiles, publicly available from the web site of the
 10 National Public Health Institute, Finland, were used. For LDL peak particle size, the cut point of 25.5 nm was used to code individuals with small LDL particles as affected. Although LDL-C is an important component trait of FCHL, serum TC was used instead in the ascertainment of the Finnish FCHL families as well as in the statistical analyses of the SNPs forming the *USF1* susceptibility haplotype. The
 15 reasoning for this is the significant hypertriglyceridemia associated with FCHL. The Friedewald formula is generally not recommended when TGs are over (400 mg/dl i.e. 4.4 mmol/l), which is often the case with hypertriglyceridemic FCHL family members. In addition, the population percentile points of LDL-C could not be estimated when including this factor, as we currently don't have population
 20 percentiles for LDL-C.

BIOCHEMICAL ANALYSES

Serum lipid parameters and LDL peak particle size were measured as described earlier^{4,9,39}. Probands or hyperlipidemic relatives who used lipid-lowering drugs were
 25 studied after their treatment was withheld for 4 weeks. In the 60 FCHL families, DNA and lipid measurements were available for 721 and 771 family members, respectively. In these 60 FCHL families, there were 226 individuals with TC > 90% age-sex specific Finnish population percentile, 220 with TGs > 90% age-sex specific percentile, 321 with TC and/or TGs > 90% age-sex specific percentile; and 125
 30 individuals with both TC and TGs > 90% age-sex specific percentiles, respectively. A total of 96 men and 124 women exhibited high TGs (>age-sex 90th percentile).

SEQUENCING, GENOTYPING AND SEQUENCE ANNOTATIONS

The *TXNIP* gene was sequenced in the 60 FCHL probands and the *APOA2*, *RXRG*,
 35 and *USF1* genes in the 31 probands of the original linkage study⁴. For *TXNIP* and *USF1*, 2000 bp upstream from the 5' end of the gene were also sequenced. For

5 *USF1*, the DNA binding domain was also sequenced in the remaining 29 probands. For all genes, both exons and introns were sequenced, except for the large 44,261-bp *RXRG* gene where only exons and 100 bp exon-intron boundaries were sequenced. Sequencing was done in both directions to identify heterozygotes reliably. Sequencing was performed according to the Big Dye Terminator Cycle

10 Sequencing protocol (Applied Biosystems), with minor modifications and the samples separated with the automated DNA sequencer ABI 377XL (Applied Biosystems). Sequence contigs were assembled through use of Sequencher software (GeneCodes). The dbSNP and CELERA databases were used to select SNPs. Pyrosequencing and solid-phase minisequencing techniques were applied

15 for SNP genotyping, as described earlier^{4,40}. Pyrosequencing was performed using the PSQ96 instrument and the SNP Reagent kit (Pyrosequencing AB). Every SNP was first genotyped in a subset of 46 family members from 18 of the 60 FCHL families. If the SNP was polymorphic (minor allele frequency > 10% in this subset), the SNP was genotyped in 238 family members of 42 FCHL families, including the

20 31 FCHL families of the original linkage study⁴. This strategy was not applied for the *TXNIP* gene the variants of which all had a minor allele frequency <10%. The physical order of the markers and genes was determined using the UCSC Genome Browser. The novel SNPs characterized in this study will be submitted to public databases (NCBI). All SNPs were tested for possible violation of Hardy Weinberg

25 equilibrium (HWE) in three groups (all family members, probands, and spouses) using the HWSNP program developed by Dr. Markus Perola at the National Public Health Institute of Finland. Annotation data of the Alu elements were downloaded from the UCSC Genome Browser, which uses the RepeatMasker to screen DNA sequences for interspersed repeats. The positions of the 60-bp sequence on these

30 Alu elements were identified using the BLAST. Other annotation data were downloaded from the LocusLink.

EXPRESSION ARRAY ANALYSIS OF ADIPOSE TISSUE

35 Six affected FCHL family members exhibiting the susceptibility haplotype (see Results) and four affected FCHL family members homozygous for the protective haplotype were selected for assessment of gene expression. All six susceptibility

haplotype carriers were from six individual families. The four homozygous protective haplotype carriers were two sibpairs from two families. Biopsies were taken from umbilical subcutaneous adipose tissue under local anaesthesia to collect 50-2000 mg of adipose tissue. The RNA was extracted using STAT RNA-60 reagent (Tel-Test, Inc.), according to the manufacturer's instructions, followed by DNase I treatment and additional purification with RNeasy Mini Kit columns (Qiagen). The quality of the RNA was assessed using the RNA 6000 Nano assay in the Bioanalyzer (Agilent) monitoring for ribosomal S28/S18 RNA ratio and signs of degradation. The concentration and the A260/A280 ratio of the samples were measured using a spectrophotometer, the acceptable ratio being 1.8-2.2. Then 2 μ g of total RNA was reverse transcribed to cDNA using the SuperScript Choice System (Invitrogen) and T7-oligo(dT)₂₄ primer, according to instructions provided by Affymetrix, except using 60 pmols of primer and a reaction volume of 10 μ l, after which biotin-labeled cRNA was created using Enzo® BioArray™ HighYield™ RNA Transcript Labeling Kit (Affymetrix). Prior to hybridization the cRNA was fragmented to obtain a transcript size distribution of 50 to 200 bases, after which samples were hybridized to Affymetrix Human Genome U133A arrays and scanned in accordance with the manufacturers' recommendations.

Scanned images were analyzed with Affymetrix Microarray Suite 5 (Affymetrix, Santa Clara, CA) software employing the Statistical Expression Algorithm. All analysis parameters were set to the default values recommended by Affymetrix. Global scaling to a target intensity of 100 was applied to all arrays but no further normalizations were performed at this point. Output files of result metrics, including the scaled signal intensity values and the corresponding detection call expressed as absent, marginal or present, were further processed using GeneSpring 5.0 data analysis software (Silicon Genetics, Redwood City, CA). For each probe array a per gene normalization was applied so that signal intensities were divided by the median intensity calculated using all 10 probe arrays. Cut-off values to discriminate low quality data were determined separately for each haplotype group by dividing the base value with the proportional value estimated using the Cross Gene Error Model implemented in GeneSpring. To identify differentially expressed genes between the two haplotypes, ratios of averaged normalized intensities were

5 calculated. Differences were considered as significant if the resulting ratio fell at
 least three standard deviations outside the average ratio calculated from the
 distribution of the \log_{10} of the ratios. To further increase result stringency only genes
 scored as present in all 10 samples, or as absent or marginal in all cases and
 present in all the controls (or vice versa), were included. Annotation information
 10 defining the biological processes that each gene could be ascribed to was retrieved
 from the classifications provided by the gene ontology (GO) consortium⁴¹. Statistical
 evaluation of enrichment of categories represented in each gene list, compared to
 the proportion observed in the total population of genes on the probe array, was
 performed using the Expression Analysis Systematic Explorer (EASE) tool⁴¹, with
 15 the threshold value set to 3. The test statistic was calculated using Fisher's exact
 test. To maximize robustness, an EASE score (p-value) was calculated where the
 Fisher exact probabilities were adjusted so that categories supported by few genes
 were strongly penalized, while categories supported by many genes were negligibly
 penalized. EASE scores (p-values) falling below 0.05 were considered statistically
 20 significant.

QUANTITATIVE REAL-TIME PCR ANALYSIS OF *USF1*

Two affected FCHL family members exhibiting the susceptibility haplotype and two
 affected FCHL family members without the haplotype were selected for assessment
 25 of *USF1* expression in adipose tissue utilizing the SYBR-Green assay (Applied
 Biosystems). Two step RT-PCR was done using TaqMan Gold RT-PCR kit
 according to manufacturers' recommendations. A total of 1 μ g of RNA was
 converted to cDNA in a 100 μ l reaction of which 1 μ l was used in the quantitative
 PCR reaction. The ratio of *USF1* to two housekeeping genes GAPDH and HPBGD
 30 was used to normalize the data. The specificity of the reaction was evaluated using
 a dissociation curve in addition to a no-template control. The following PCR primers
 were used in separate 10 μ l SYBR-Green reactions: For *USF1*; forward: 5'-
 ATGACGTGCTTCGACAACAG-3', reverse: 5'-GGGCTATCTGCAGTTCTTGG-3'.
 For GAPDH; forward: 5'-CGGAGTCAACGGATTTGGTCGTAT3', reverse: 5'-
 35 AGCCTTCTCCATGGTGGTGAAGAC-3'. For HPBGD; forward: 5'-
 AACCTCATGATGCTGTTGTC-3', reverse: 5'-TAGGATGATGGCACTGAACTC3'.

- 5 The reactions were run in triplicate using the ABI Prism 7900 HT Sequence Detection System in accordance with the manufacturers' recommendations and the data were analyzed using Sequence Detector version 2.0 software.

INITIAL FUNCTIONAL ANALYSIS

- 10 Initial functional analyses were performed using the SEAP reporter system (Clontech Laboratories, Palo Alto, CA) in COS cells. This system utilizes SEAP, a secreted form of human placental alkaline phosphatase, as a reporter molecule to monitor the activity of potential promoter and enhancer sequences. The constructs were cloned into the pSEAP2-Enhancer vector which contains the SV40 enhancer.
- 15 The correct allele and orientation in each construct was verified by sequencing. Cell culture media between 48 h and 72 h after transfection were taken for the SEAP reporter assay. The monitoring of the SEAP protein was performed using the fluorescent substrate 4-methylumbelliferyl phosphate (MUP) in a fluorescent assay according to the manufacturer's instructions. Data are representative of at least two
- 20 independent experiments.

STATISTICAL ANALYSES

- Parametric linkage and nonparametric affected sib-pair (ASP) analyses were carried using the same programs and parameters as in the original linkage study⁴. Two
- 25 traits were investigated, the FCHL and TG trait. The MLINK program of the LINKAGE package⁴³ version FASTLINK 4.1P⁴⁴⁻⁴⁵ was used as implemented by the ANALYZE package⁴⁶ to perform the parametric two-point and multipoint linkage analyses. The ASP analysis was performed using the SIBPAIR program of the ANALYZE package⁴⁶. For each marker, allele frequencies were estimated from all
- 30 individuals using the DOWNFREQ program⁴⁷.

- The SNPs were tested for association using the HHRR²⁷ and the gamete competition test²⁹. To minimize the number of tests performed, the SNPs residing outside the USF1-JAM1 region were tested for association only using the HHRR²⁷ test when analyzing the TG- and FCHL-affected males. The HHRR analysis,
- 35 performed by use of the HRRLAMB program⁴⁸, tests the homogeneity of marker allele distributions between transmitted and non-transmitted alleles. The multi-

5 HHRR analysis is testing the same hypothesis using several SNPs. The gamete competition test is a generalization of the TDT and views transmission of marker alleles to affected children as a contest between the alleles, making effective use of full pedigree data. The gamete competition method is not purely a test of association, because the null hypothesis is no association and no linkage, and thus linkage in itself also affects the observed p-value. Furthermore, the gamete competition test readily extends to two linked markers, enabling simultaneous analysis of multiple SNPs in a gene. P-values based on asymptotic approximations can be biased when data used to calculate them are relatively sparse. To confirm that the gamete competition results are indeed significant we also calculated empirical p-values for all analyses involving multiple SNPs (Table 1) using gene dropping. In gene dropping the founder genotypes are assigned using the estimated allele frequencies assuming HWE and linkage equilibrium (LE). The offspring genotypes are assigned assuming Mendelian segregation. Thus gene dropping is performed under the null hypothesis of LE and no linkage. To calculate an empirical p-value, gene dropping is performed multiple times. Here at least 50,000 simulations were performed for each analysis. The likelihood ratio test statistic (LRT) from each gene dropping iteration is compared to the LRT for the observed data. The empirical p-value is the proportion of iterations in which the gene dropping LRT equaled or exceeded the observed LRT. In general, the obtained empirical p-values of gene dropping are more conservative than asymptotic p-values for small sample sizes.

The HBAT program, options optimize offset (-o) and empirical test (-e), were performed to test for association between haplotypes and the trait⁴⁹. The option -o measures not only preferential transmission of the susceptibility haplotype to affecteds but also less preferential transmissions to unaffecteds. The -e option leads to a test of association given linkage and gives thus an empirical estimation of the variance. These haplotype analyses are affected by the fact that four of the 15 SNPs for the *JAM1-USF1* region were genotyped in the 60 extended FCHL families and 11 SNPs in 42 nuclear FCHL families. The genotype Pedigree Disequilibrium Test (geno-PDT)⁵⁰, which provides a genotype-based association test for general pedigrees, was also performed for a combination of genotypes from selected *USF1*

5 SNPs (Table 3). LD between the marker genotypes for SNPs in the *JAM1-USF1*
region was tested using the Genepop v3.1b program, option 2, at their web site. In
this program, one test of association is performed for genotypic LD, and the null
hypothesis is that genotypes at one locus are independent from the genotypes at
the other locus. The program creates contingency tables for all pairs of loci in each
10 population and performs Fisher exact test for each table using a Markov chain.

URLs

Supplementary Tables 1-4 and further details on microarray data will be available at
our web site (www.genetics.ucla.edu/labs/pajukanta/fchl/chr1/). The raw data for the
15 complete set of probe arrays can be accessed through the Gene Expression
Omnibus at NCBI (www.ncbi.nlm.nih.gov/geo) using the GEO accession GSE590.
The Finnish 90th age-sex specific percentile values for TC and TGs are available at
the web site of the National Public Health Institute of Finland
(www.ktl.fi/molbio/wwwpub/fchl/genomescan). We used the dbSNP (available at
20 www.ncbi.nlm.nih.gov) and CELERA (www.celera.com) for SNP selection; the
UCSC Genome Browser (genome.ucsc.edu) for physical order of the genes and for
annotation of the Alu element; the BLAST (www.ncbi.nlm.nih.gov/blast/) for blasting
sequences against human and mouse databases; the LocusLink
(www.ncbi.nlm.nih.gov/LocusLink/) to download annotation data; and the Genepop
25 (wbiomed.curtin.edu.au/genepop/index.html) to calculate intermarker LD.

CLAIMS

5

1. A nucleic acid molecule comprising a chromosomal region contributing to or indicative of hyperlipidemias and/or dyslipidemias and/or defective carbohydrate metabolism, wherein said nucleic acid molecule is selected from the group consisting of:

10

(a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence has one or more mutations having an effect on USF1 function;

15

(b) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence is characterized by comprising a guanine or an adenine residue in position 3966 in intron 7 of the USF1 sequence; and/or

20

(c) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence is characterized by comprising a cytosine or a thymine residue in position 5205 in exon 11 of the USF1 sequence;

wherein said nucleic molecule extends, at a maximum, 50000 nucleotides over the 5' and/or 3' end of the nucleic acid molecule of SEQ ID NO: 1.

2. The nucleic acid molecule of claim 1 which is genomic DNA.

25

3. A fragment of the nucleic acid molecule of claim 1 or 2 having at least 20 nucleotides wherein said fragment comprises nucleotide position 3966 and/or position 5205 of SEQ ID NO:1.

4. A nucleic acid molecule which is complementary to the nucleic acid molecule of any one of claims 1 to 3 and which has a length of at least 20 nucleotides.

5. A vector comprising the nucleic acid molecule of any one of claim 1 to 4.

- 5 6. A primer or primer pair, wherein the primer or primer pair hybridizes under stringent conditions to the nucleic acid molecule of any one of claims 1 to 4 comprising nucleotide positions 3966 and 5205 SEQ ID NO:1 or to the complementary strand thereof.
7. A non-human host transformed with the vector of claim 5.
- 10 8. The non-human host of claim 7 which is a bacterium, a yeast cell, an insect cell, a fungal cell, a mammalian cell, a plant cell, a transgenic animal or a transgenic plant.
9. A pharmaceutical composition comprising USF1 or a fragment thereof, a nucleic acid molecule encoding USF1 or a fragment thereof, or an antibody specific for USF1.
- 15 10. A diagnostic composition comprising a nucleic acid molecule encoding USF1 or a fragment thereof, the nucleic acid molecule of any one of claims 1 to 4, the vector of claim 5, the primer or primer pair of claim 6 or an antibody specific for USF1.
- 20 11. A method for testing for the presence or predisposition of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism, comprising analyzing a sample obtained from a prospective patient or from a person suspected of carrying such a predisposition for the presence of a wild-type or variant allele of the USF1 gene.
- 25 12. The method of claim 11, wherein said variant comprises an SNP at position 3966 and/or at position 5205 of the USF1 gene in a homozygous or heterozygous state.
13. The method of claim 11 or 12, wherein said testing comprises hybridizing the complementary nucleic acid molecule of claim 4 under stringent conditions to nucleic acid molecules comprised in a sample and detecting said hybridization,
- 30

- 5 wherein said complementary nucleic acid molecule comprises the sequence position containing the SNP.
14. The method of any one of claim 11 to 13 further comprising digesting the product of said hybridization with a restriction endonuclease or subjecting the product of said hybridization to digestion with a restriction endonuclease and
10 analyzing the product of said digestion.
15. The method of claim 14, wherein said probe is detectably labeled.
16. The method of any one of claims 11 to 15, wherein said testing comprises determining the nucleic acid sequence of at least a portion of the nucleic acid molecule of any one of claims 1 to 4, wherein said portion comprises the
15 position of the SNP.
17. The method of claim 16, wherein the determination of the nucleic acid sequence is effected by solid-phase minisequencing.
18. The method of claim 17 further comprising, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid
20 molecule.
19. The method of claim 11 to 15, wherein said testing comprises carrying out an amplification reaction wherein at least one of the primers employed in said amplification reaction is the primer of claim 6 or belongs to the primer pair of claim 6, comprising assaying for an amplification product.
20. The method of claim 19 wherein said amplification is effected by or said
25 amplification is the polymerase chain reaction (PCR).
21. A method for testing for the presence or predisposition of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism comprising assaying a sample obtained from a human for the amount of USF1 contained
30 in said sample.

- 5 22. The method of claim 21, wherein said testing is effected by using an antibody or aptamer specific for USF1.
23. The method of claim 22, wherein said antibody or aptamer is detectably labeled.
- 10 24. The method of any one of claims 21 to 23, wherein the test is an immunoassay.
25. The method of any one of claims 11 to 24, wherein said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human tissue.
- 15 26. The method of any one of claims 11 to 25, wherein the nucleic acid molecule or protein from said sample is fixed to a solid support.
27. The method of claim 26, wherein said solid support is a chip, a silica wafer, a bead or a microtiter plate.
- 20 28. Use of the nucleic acid molecule of any one of claims 1 to 5 for the analysis of the presence or predisposition of hyperlipidemia and/or dyslipidemia and/or defective carbohydrate metabolism.
- 25 29. Use of USF1 or a fragment thereof or of a nucleic acid molecule encoding USF1 and/or comprising at least the wild-type sequence of intron 7 and/or exon 11 of USF1, for the preparation of a pharmaceutical composition for the treatment of hyperlipidemias and/or dyslipidemias including familial combined hyperlipidemia (FCHL), hypercholesterolemia, hypertriglyceridemia, hypoalphalipoproteinemia, hyperapobetalipoproteinemia (hyperapoB), familial dyslipidemic hypertension (FDH), metabolic syndrome, type 2 diabetes mellitus, coronary heart disease, atherosclerosis or hypertension.

- 5 30. Kit comprising the nucleic acid molecule of any one of claims 1 to 5, the primer or primer pair of claim 6 and/or the vector of claim 7 in one or more containers.

17 Feb. 2004

Abstract

5

10


15

20

25

The present invention relates to a nucleic acid molecule comprising a chromosomal region contributing to or indicative of hyperlipidemias and/or dyslipidemias or defective carbohydrate metabolism, wherein said nucleic acid molecule is selected from the group consisting of: (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence has one or more mutations having an effect on USF1 function; (b) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence is characterized by comprising a guanine or an adenine residue in position 3966 in intron 7 of the USF1 sequence; and/or (c) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, wherein said nucleic acid sequence is characterized by comprising a cytosine or a thymine residue in position 5205 in exon 11 of the USF1 sequence; wherein said nucleic acid molecule extends, at a maximum, 50000 nucleotides over the 5' and/or 3' end of the nucleic acid molecule of SEQ ID NO: 1. The present invention further relates to a diagnostic composition comprising a nucleic acid molecule encoding USF1 or a fragment thereof, the nucleic acid molecule disclosed herein, the vector, the primer or primer pair of the present invention or an antibody specific for USF1. Finally, the present invention relates to the use of the nucleic acid molecule of the invention for the preparation of a pharmaceutical composition for the treatment of hyperlipidemia, dyslipidemia, coronary heart disease, type II diabetes, metabolic syndrome, hypertension or atherosclerosis.

Up-regulated genes



Down-regulated genes





Figure 2



A total 91 human genes have this 60 bp sequence, similar to the mouse B1 repeat, as a part of *AluSx* on the coding strand (43 genes) or on the opposite strand (48 genes)

Figure 3a




Figure 3b

17 Feb. 2004

SEQUENCE LISTING

<110> National Public Health Institute

<120> Identification of SNPs associated with hyperlipidemia, dyslipidemia and defective carbohydrate metabolism

<130> K 1114 EP

<160> 1

<170> PatentIn version 3.1

<210> 1

<211> 5687

<212> DNA

<213> Human

<220>

<221> variation

<222> (3966)..(3966)

<223> r = adenine (a) or guanine (g):
adenine is wild-type associated; guanine is disease-associated

<220>

<221> variation

<222> (5205)..(5205)

<223> y = cytosine (c) or thymine (t):
thymine is wild-type associated; cytosine is disease-associated

<400> 1
ttgaaaattt tccttggata ggaaagggtt ggaggacctt atgggtagag aatttcctaaa 60
aatcttgccc cttttgtgtt gggattatct tattgctttg tactgtgtag ctgtttcttt 120
ctggaggcat gtctgcccag ctctttgttt ttcttgcct ctggctgggt gtcagggtcc 180

taaggcagag cttgtaggtg gattcttccc cctttgtctc ttcttcagaa ccctgttttt	240
ttttttttta ccccttcttg ctcaggctta gttgatttgg agttgtcata gcaacatttt	300
agcaacagtg ttgttctgca ggaaggcttg atgaataaaa tagagaatgc ttgaagagga	360
tccacttggg ctttaggggt tctaacagat tatataaatc tggatacccc aaaacaagag	420
tcctgtcagt agaattggggc ccaaagcca agtctagtct ttgtggtcag ggatattctt	480
ccagtggtag tgggcttcag atttctctt cctaggtttg aaaacagaaa tgtcttgatg	540
gacaacatgt ggctgagaaa ctggaagaag catcagtgtc catgacactg tttttttga	600
tgggtggggcc aatacatggc ccttctgat tcccatgaag ctgccatcat ggcaggtc	660
aatagcttta atgatccatt tagagatgtg ttgttggctg ggtgcggtgg ctcatgcctg	720
taatccaagc actttgggag gccgaggcag gccgatcacc tgaggtcagg agttccagac	780
cagcctggcc aatatggtaa aaccccatct ctactgaaaa taaaaaatt agctgggcgt	840
ggtggtgggc acctataatc ccagctattc aggaggctga ggcaggagaa tcacttgaac	900
ccaggagatg gaggttgtaa gccgagattg tgccactgca ctccagcctg ggtgacagag	960
caagattctg tctcagaaaa aaaaaaaaaa aaaagaaaga aatgtgttgt ttcggccagg	1020
tgcagtggct cacacctgta atcccagcac tttgggaggc tgccgagggtg gacagatcat	1080
gctctcagga gtctgagacc agccgggcca acatggtgaa acccgtctc tactaaaaat	1140
acaaaaatta gccaggcgtg gtggtgtgca cctgtaatcc cagctactcc ggaggctgag	1200
gcaggagaat cacttgaacc tgggaggcag aggttgcagt gagctgagat cgcgccactg	1260
cactccagcc tgggtgacag agagagactc tgtctcaaaa aaaaaaaaaa aaaaaagtg	1320
ttgttctgt cttccagtat aattatccac tctccaccag gagttggagt gataatggag	1380
ggatggggaa cactatttgt agccttgctt tttcaatcac tgtaggccag tctcaacat	1440
cagtatggtg gaggtgatt gtcccctgca gatgactggg ttattttcct ggctatgtgt	1500
tcattggaacc taagttctag aaccagagat actgttctgt ttctaaact cattgcaaac	1560
ttcatgattt ctaccaggac ttagcactca ggcctgtgaa tcaggagata caaagacctc	1620
caaaaaagga ccagttcctc ggatgtgccc cctcacagag agatgaagggt gtgagtgaag	1680
aagaggtagg gtctgggatg aaagatgggt ggcctggaag aatgcaaaat gaccaagagc	1740
actgcctctg gagtcaggca gacctggatt caggttctac tctatcactt actgtgtgat	1800
ttggtttctc tatctataaa atggaagtag tgctatctat ctctgtgtgc tgtttttagt	1860
actaaataag attacatgta atgtacttag cttagtgtt atgtacatag taaacagtaa	1920
acactagtgt ttattctaac ctaaccagc ttctgttggg aatgccaatg agtttgcagc	1980
catatgttac tgggccagtg agcttctcat tgacttcttc tcatactctt ccttttgtcc	2040
tttcaccaca aacaggcagc agaaaacagc tgaaacggaa gaggggacag tgcagattca	2100

ggaaggtgag	tgctagaaac	agaaccaaga	ctaagaaccc	atcatggcct	cccttccttc	2160
cccaccagac	catctcctgt	gcatectect	ccttcctgtga	catgcaaagt	gaacgggggt	2220
agaaaggcag	ttaaactcaca	gacttttctt	ttgttctttt	aattcaggtg	cagtggctac	2280
tggggaagac	ccaaccagtg	tggctattgc	cagcatccag	tcagctgcca	ccttccttga	2340
ccccaacgtc	aagtacgtct	tccgaactga	gaatgggggc	caggtaaggg	agggggccag	2400
gtggctgcag	gtgttatctg	gggttgggat	tgaggaggt	aattgaacat	gtcttgggga	2460
gacctggcct	ggaggatgag	ttgaaagagt	ggactgttgc	aggggagggg	ggtgctaata	2520
ctggagtaga	gactgggtgtg	aggtagatg	tatgctgaaa	cctctgtgtg	gggaaagaag	2580
ggagaatggc	tgaatccatg	tctctgaagg	actttgtttt	ggggccctat	ccaagggaag	2640
ctttatgagg	ggccctagga	ttcccaacac	ttaatctttt	cttctctctt	cactccctct	2700
gccttcctct	acacttctag	gtgatgtaca	gggtgatcca	ggtgtctgag	gggcagctgg	2760
atggccaaac	tgagggaact	ggcgccatca	gtggctaccc	tgccactcaa	tccatgaccc	2820
aggtagcagg	tatgggctgg	ggaggtcact	agagttctga	gaagtaagat	gaagaaggga	2880
atcagtagga	tgggggtgaa	gctaggaaca	gtgaggcatc	taaggctgcc	ttgtcccaaa	2940
gcactaggct	ctccttttct	ggatgtttct	ctctctctct	ctctctctct	ccaccctacc	3000
taccacccca	acggatagaa	gctgcagagt	ggtgtagtgg	gaagaagttt	ttgactgtta	3060
ccagaatcag	ttttcttgct	ccccttccca	ggcgggtgatc	cagggtgctt	tcaccagtga	3120
tgatgcagtt	gacacggagg	ggacagctgc	tgagacgcac	tatacttact	tccccagcac	3180
ggcagtggga	gatggggcag	gggttaccac	atcggggagt	acagctgctg	ttgttactac	3240
ccagggtcga	gaggcactgc	tggggcaggc	gacccctcct	ggcactgggtg	agatattgca	3300
tgaggatgct	ggctgaaagg	gctagaatag	gctgtgggac	atgactggta	ggcagtgagc	3360
cttcactcat	gactcttagt	gatcattaag	acctggacag	gcagtgagtc	tggggctgct	3420
cttctattag	catgttcttt	ttagaggagg	ggaccagggg	cttcacctca	gggcttggtg	3480
aggttectac	ccatgtcctg	acagaacctc	ccctgcctct	tcacaggtca	attctttgtg	3540
atgatgtcac	cacaagaagt	actgcaggga	ggaagccagc	gctcaattgc	ccctaggact	3600
cacccttatt	ccccgtgagt	gacccttggt	tcttctcaga	ttccgtaagt	ggtttttttt	3660
tttttttttt	ttttttgaga	cagagtcttg	ctctgtcacc	caggctggag	tgcagtggca	3720
tgatctcagc	tcaactgcaac	ctctgcttcc	agggttcaag	cgtttctcat	gcctcagcct	3780
cctgagtagc	tggaaactaca	gacatgtacc	accacccctg	gctaattttt	gtatctttag	3840
tagagacagg	gtttcaccat	gttggccagg	ctgggtctga	actcctgacc	tcaagtgate	3900
cgctgcctc	ggcctcccaa	agtgtctggg	ttacaggtgt	gagacaccac	acctagctac	3960
cataartggg	cctaatacct	gctaaatctt	gtataattcc	ttaaccccaa	acttcaatca	4020

tgtatTTTTgt	cttcttactc	tggccaccct	gggctctgtt	gtcaggaagt	cagaagctcc	4080
ccggacgact	cgggatgaga	aacgcagggc	tcagcataat	gaaggtaggt	atgatctggg	4140
tggagctaga	agctgtctgg	tgtgatctca	gcagtgatgt	ctgaggggag	gagggattag	4200
gtaatTTTtac	cctgggactt	gtggcgagtt	ttcactgagt	caccttgtcc	tccactttgc	4260
cccacagtgg	agcgtcgccg	ccgagacaag	atcaacaact	ggatcgtgca	gctctccaag	4320
ataatcccag	actgctctat	ggagagcacc	aagtctggcc	aggtcatgga	aagaccctgg	4380
tagtgggcag	gatgcctgaa	ttctgcctcc	tggatattgtt	tccagaaatg	gtagagagag	4440
gggcacacat	gacagtagtc	ttatctctcc	ctgaggttcc	tgtatccctg	ggagatatta	4500
taccaccttc	cttagatgaa	aatgaggctc	aaagtgtgaa	cctacttttg	gaaagcaagc	4560
tgggtatctg	aaatcctagt	tctcatTTtg	ttgaccttat	cttgacagagt	aaaggTggga	4620
ttctatccaa	agcttTgtgat	tatatccagg	agcttccggca	gagtaaccac	cgcttTgtctg	4680
aagaactgca	gggacttgac	caactgcagc	tggacaatga	cgtgcttcga	caacaggTca	4740
gactcctacc	cccagtgcag	cccttctcag	ttctgctagc	cactgaccca	gtttgacacc	4800
ctctactttg	ttctccatgg	agaaggcttc	atcttttccc	cctcaccagt	ggatgtctga	4860
atacattcag	gggcttggaa	gtgccagctt	tactacctat	tccctttact	gcctccttcc	4920
catgtcaggt	ggaagatctt	aaaaacaaga	atctgctgct	tcgagctcag	ttgcggcacc	4980
acggattaga	ggtcgtcatc	aagaatgaca	gcaactaact	atggggattc	aggggctttg	5040
ggcccaagaa	ctgcagatag	cccaggagca	acagcctaata	cccgTgcccc	tttcccttcac	5100
tgccccactt	ctggcatggg	acagggggaa	gttcagaagg	tgtgtccttg	aactgaggcc	5160
ctgtgatatg	gcggcctgca	gtggtgtgaa	acacacaatg	tggaygtgca	ctgacagcct	5220
tgccccaccc	caccatgcag	cccctggggc	cttgtgctcc	tctcgacaaa	tgcattgtgt	5280
gtctccatgc	tggatactgg	acacactaaa	ctctggggct	tgtcctgtgc	ttgcttagag	5340
tgcccagcag	aggtttTgtg	acaggTgatg	ctctggcttg	ccccaggact	ctggcacttc	5400
cattggTtct	tcctttccct	ggagctgagg	tttagatgtg	caacctgtgg	ctcaggggag	5460
caagcttaca	caagaagtga	gggaaggatg	tttagcagtg	gctggTgccc	atgaagagga	5520
gattggccag	tgagaagctg	aggcctatgc	agacatctct	ggagccagag	agaacaacag	5580
gcagggggccc	acttggggcc	ttcccccttg	tgggggtcgt	TTTTTTTTTT	tcttttcttt	5640
TTTTTTTTTT	TTTTTTTTTT	TTTTtaagat	aaaattgttc	aaagcca		5687